

# **Stabilitas isoniazid dalam sediaan extemporaneous = Stability of isoniazid in extemporaneous preparations**

Monica Sari Dewi, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20510771&lokasi=lokal>

---

## **Abstrak**

Tuberkulosis merupakan salah satu penyebab kematian utama di dunia dan di Indonesia yang diakibatkan oleh infeksi bakteri *Mycobacterium tuberculosis*. Pengobatan tuberkulosis merupakan pengobatan jangka panjang dengan memberikan obat dalam bentuk kombinasi dari beberapa jenis obat antibiotik. Salah satu obat yang paling umum digunakan adalah Isoniazid. Sediaan Isoniazid yang beredar secara komersial umumnya tersedia dalam bentuk tablet. Pasien pediatri umumnya mengalami masalah dari segi farmakoterapi obat yaitu kesulitan dalam menelan obat, terutama obat sediaan padat.

Peracikan obat dalam bentuk serbuk menjadi solusi utama dalam mengatasi hal tersebut, namun obat racikan serbuk menimbulkan permasalahan di praktik kefarmasian. Sediaan racikan *extemporaneous* dibuat untuk menggantikan obat racikan serbuk karena dinilai memiliki stabilitas yang lebih baik dan dapat menutupi rasa pahit dari obat dalam bentuk racikan serbuk sehingga cocok untuk pasien pediatri. Dalam review artikel ini akan dibahas mengenai stabilitas isoniazid dalam sediaan oral cair *extemporaneous* untuk pasien pediatri, dan faktor yang mempengaruhi stabilitas sediaan *extemporaneous*.

.....

Tuberculosis is one of the major causes of death in the world and in Indonesia that comes from an infection by *Mycobacterium tuberculosis*. The treatment for curing Tuberculosis is a long-term treatment by giving drugs in combination of several types of antibiotic. One of the most commonly Tuberculosis drugs is Isoniazid. Isoniazid commercially available in tablet form. Some patients, especially the pediatric have problems in swallowing drugs that were formulated in solid or tablet form.

Compounded drug in powder form becomes the main solution to resolve this problem, however the powder concoction drugs has some problems in pharmaceutical practice. Those problems made the pharmacists began to formulate extemporaneous preparation to replace the powder concoction because extemporaneous formulation considered to have better stability and could mask the bitter aftertaste of drugs in powder concoction, making its suitable for pediatric.

This article review highlights the stability of isoniazid in extemporaneous oral liquid preparations for pediatric, also the main factors of the stability in extemporaneous preparations.