

# **Formulasi Sediaan Sirup Ekstrak Daun Kelor (*Moringa oleifera Lam.*) Sebagai Antianemia = Formulation of Moringa Leaf Extract Syrup (*Moringa oleifera Lam.*) As Antianemia**

Faiqoh Zaqladi, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20513427&lokasi=lokal>

---

## **Abstrak**

Anemia merupakan salah satu masalah kesehatan masyarakat yang mempengaruhi penduduk di negara maju maupun berkembang. Penyebab tertinggi anemia adalah kekurangan zat besi. Anemia Defisiensi Besi (ADB) dapat menyebabkan gangguan perkembangan perilaku, kognitif, dan keterampilan psikomotori anak. Daun kelor dapat menjadi alternatif potensial dalam memenuhi kebutuhan zat besi karena memiliki kandungan zat besi 9 kali lebih banyak daripada bayam. Suplementasi bubuk daun kelor terbukti dapat menurunkan prevalensi anemia sedang dan berat berturut-turut sebesar 68,2% dan 77,9% pada anak-anak berusia di bawah 2 tahun. Pemanfaatan daun kelor sebagai tanaman obat yang memiliki berbagai manfaat bagi kesehatan umumnya masih secara tradisional. Rasa pahit pada kelor menyebabkan anak-anak tidak menyukai daun kelor. Tujuan dari penelitian ini adalah mendapatkan formula sediaan sirup ekstrak daun kelor yang memiliki stabilitas fisik dan kimia yang baik. Ekstrak daun kelor diperoleh dengan Microwave Assisted Extraction (MAE). Formula sirup dibuat dengan 3 konsentrasi propilen glikol sebagai kosolven dan pengawet. Hasil yang diperoleh menunjukkan bahwa formula 1 memiliki stabilitas fisik yang baik, dan stabilitas kimia yang paling baik dengan kandungan zat besi yang paling tinggi yaitu 2,83 mg / 30 gram ekstrak / 100 ml sediaan. Kadar zat besi yang disarankan untuk anak adalah 8-10 mg/hari sehingga ekstrak yang dibutuhkan yaitu 84,80-106,00 gram ekstrak / hari dan dosis sediaan sirup yang dibutuhkan yaitu 283-353 ml/hari.

.....Anemia is one of the health problems that affect people in both developed and developing countries. The highest cause of anemia is iron deficiency. Iron Deficiency Anemia (IDA) can cause impaired development in behaviour, cognition, and psychomotor skills of children. Moringa leaves can be a potential alternative in meeting iron needs because they contain 9 times more iron content than spinach. *Moringa oleifera Lam.* leaf powder supplementation can reduce the prevalence of moderate and severe anemia in the by 68.2% and 77.9% in children below two years. The use of *Moringa oleifera Lam.* leaves as a medicinal plant with health benefits is still generally traditional. The bitter taste of *Moringa oleifera Lam.* leaves causes children to dislike *Moringa oleifera Lam* leaves. The purpose of this work was to obtain syrup formula for antianemia using Moringa leave's extract that has physical and chemical stability. Moringa leave's extract was obtained with Microwave-Assisted Extraction (MAE). Syrup formula was made in 3 concentration of propylene glycol as a cosolvent and preservatives. Result of this study showed that formula 1 has good physical stability, and the best chemical stability with the highest iron content, 2.83 mg/30 gram extract/100 ml. The recommended iron level for children is 8-10 mg / day, so the required extract is 84.80-106.00 grams of extract / day and the required dosage of syrup is 283-353 ml/day