

# Efektifitas Positioning Lateral Kanan Kiri 30° dan Positioning Lateral Kanan Kiri 90° Terhadap Keberhasilan Weaning Di Ruang Perawatan Intensif = The Effectiveness of Left Left Lateral Positioning 30A° and Right Left Lateral Positioning 90A° Against Weaning Success In the Intensive Care Room

Andi Kurniawan, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20520301&lokasi=lokal>

---

## Abstrak

Positioning yang optimal pada pasien akan berdampak pada airway manajemen dan ventilasi, menjaga keselarasan posisi tubuh, serta memberikan keamanan dan kenyamanan fisiologis. Tindakan positioning sudah dilakukan di ICU RSUD Prof Dr. Margono Soekarjo dengan tujuan untuk mengurangi resiko luka dekubitus, tetapi belum dilihat efeknya pada kemampuan fisiologis paru pasien yang terpasang ventilator. Dari berbagai bentuk tindakan positioning yang dapat dilakukan terhadap pasien hingga saat ini belum ada protokol – protokol khusus yang menjelaskan terkait positioning mana yang paling efektif dan efisien terhadap pasien khususnya pada pasien dengan kejadian prolong ventilator diruang ICU. Tujuan dari penelitian ini yaitu mengetahui pengaruh dan efektifitas positioning lateral kanan kiri 30 ° dan positioning lateral kanan kiri 90 ° terhadap keberhasilan weaning pada pasien dengan prolong ventilator mekanik di ruang ICU. Sampel dalam penelitian ini berjumlah 40 responden yang dibagi menjadi 20 responden kelompok intervensi 1 ( positioning lateral kanan kiri 90 ° ) dan 20 responden kelompok intervensi 2 ( positioning lateral kanan kiri 30 ° ). Metode penelitian ini adalah quasy eksperiment dengan desain Pre test and post test pada kedua kelompok intervensi ( with control group ). Hasil penelitian menunjukan positioning lateral kanan kiri 30 ° berpengaruh secara sigifikan terhadap keberhasilan weaning pada pasien dengan prolong ventilator mekanik ( p value 0,000 ) dibandingkan positioning lateral kanan kiri 90 ° ( p value 0,180 ). Pengaturan sudut kemiringan pada saat pemberian positioning lateral akan mengurangi tahanan otot - otot perut sehingga mengurangi tekanan ke diafragma dan meringankan kompresi dada. Hal ini akan berdampak pada masuknya udara lebih banyak ke dalam paru – paru serta meningkatnya oksigenasi dalam meningkatkan fungsi fisiologis paru. Hasil dari penelitian ini diharapkan menjadi suatu gold standar baru berupa pemberian intervensi positioning lateral kanan kiri 30 ° khususnya pada pasien dengan penggunaan ventilator diruang ICU.

.....Optimal positioning of the patient will have an impact on airway management and ventilation, maintain body position alignment, and provide physiological safety and comfort. Positioning action has been carried out in the ICU RSUD Prof. Dr. Margono Soekarjo with the aim of reducing the risk of pressure sores, but the effect has not been seen on the physiological abilities of the lungs of patients on ventilators. Of the various forms of positioning actions that can be performed on patients, there are currently no specific protocols that explain which positioning is most effective and efficient for patients, especially in patients with prolong ventilator events in the ICU. The purpose of this study was to determine the effect and effectiveness of right and left lateral positioning of 30 ° and right and left lateral positioning of 90 ° on the success of weaning in patients with mechanical ventilator prolong in the ICU. The sample in this study amounted to 40 respondents who were divided into 20 respondents in the intervention group 1 (right and left lateral positioning 90 °) and 20 respondents in the intervention group 2 (right and left lateral positioning 30 °).

°). This research method is a quasi-experimental design with pre-test and post-test in both intervention groups (with control group). The results showed that right and left lateral positioning of 30 ° had a significant effect on the success of weaning in patients with mechanical ventilator prolong (p value 0.000) compared to right and left lateral positioning of 90 ° (p value 0.180). Adjusting the angle of inclination at the time of lateral positioning will reduce the resistance of the abdominal muscles thereby reducing pressure on the diaphragm and relieve chest compressions. This will have an impact on the entry of more air into the lungs and increased oxygenation in improving the physiological function of the lungs. The results of this study are expected to become a new gold standard in the form of giving an intervention of 30° right and left lateral positioning, especially in patients using ventilators in the ICU.