

# Analisis Kelimpahan Mikroplastik pada Air, Sedimen, dan Ikan Cere *Gambusia affinis* (Baird & Girard, 1853) di Situ Kenanga dan Situ Mahoni, Universitas Indonesia, Depok, Jawa Barat = Analysis of Microplastics Abundance in Water, Sediment, and Mosquitofish *Gambusia affinis* (Baird & Girard, 1853) in Situ Kenanga and Situ Mahoni, Universitas Indonesia, Depok, West Java

Ritonga, Sabrina Agistia, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20521618&lokasi=lokal>

---

## Abstrak

Mikroplastik diketahui telah tertelan oleh berbagai taksa biota. Penelitian ini dilakukan untuk menganalisis kelimpahan, bentuk, dan warna mikroplastik yang terdapat pada air, sedimen, dan ikan cere *Gambusia affinis* di Situ Kenanga dan Situ Mahoni, Universitas Indonesia, Depok, Jawa Barat. Seluruh sampel diambil di 3 stasiun yaitu inlet, midlet, dan outlet. Sampel air 20 L disaring menggunakan plankton net 350 mesh, sampel sedimen diambil 250 mL menggunakan Ekman grab, ikan cere diambil secara acak menggunakan dip net sebanyak 20 individu dari tiap situ. Sampel ikan diawetkan dengan alkohol 70% dan diekstraksi menggunakan 10 mL NaOH 1 M. Sedimen dikeringkan pada oven 65°C dan dihaluskan. Seluruh sampel dilarutkan dengan NaCl jenuh. Larutan dihomogenisasi sebanyak 20 mL dan 1 mL ditetaskan Sedgewick Rafter Chamber untuk diamati di bawah mikroskop cahaya. Hasil pengamatan menunjukkan total rata-rata kelimpahan mikroplastik di Situ Kenanga pada air sejumlah  $434,33 \pm 23,51$  partikel L-1, pada sedimen sejumlah  $45.837,04 \pm 36.305,97$  partikel kg-1, pada ikan cere sejumlah  $287,67 \pm 95,65$  partikel ind-1 atau  $212,28 \pm 61,88$  partikel gram-1. Sementara itu, pada air di Situ Mahoni sejumlah  $437,67 \pm 30,21$  partikel L-1, pada sedimen sejumlah  $36.237,04 \pm 16.702,59$  partikel kg-1, dan pada ikan cere sejumlah  $275 \pm 95,82$  partikel ind-1 atau  $654,28 \pm 279,71$  partikel gram-1. Hasil analisis Uji T Dua Sampel menunjukkan tidak terdapat perbedaan nyata pada sampel air, sedimen, dan ikan cere (partikel ind-1), tetapi terdapat perbedaan nyata pada ikan cere (partikel g-1) dari kedua situ. Fiber dan warna hitam merupakan bentuk dan warna mikroplastik yang mendominasi keseluruhan sampel.

.....Microplastics have been ingested by various biota taxa. This study was aimed to analyze the abundance, shape, and color of microplastic found in water, sediment, and mosquitofish *Gambusia affinis* in Situ Kenanga and Situ Mahoni, University of Indonesia, Depok, West Java. All samples were taken at 3 stations, namely inlet, midlet, and outlet. Water samples (20 L) were filtered using a 350 mesh plankton net, sediment samples (250 mL) were taken using an Ekman grab, mosquitofish were taken randomly using a dip net, and 20 individuals from each situ (total of 40 individuals). Fish samples were preserved in 70% alcohol and extracted using 10 mL of 1 M NaOH. Sediments were dried in an oven at 65°C and pulverized. All samples were dissolved with saturated NaCl. The solution (20 mL) was homogenized, then 1 mL of solution was taken to the Sedgewick Rafter Chamber and then observed under a light microscope. The results showed that the average total of microplastics in Situ Kenanga in water was  $434,33 \pm 23,51$  L-1 particles, in sediments were  $45.837,04 \pm 36.305,97$  kg-1 particles, and in fish were  $287,67 \pm 95,65$  ind-1 particles or  $212,28 \pm 61,88$  gram-1 particles. Meanwhile, the average total of microplastics in Situ Mahoni in water were  $437,67 \pm 30,21$  L-1 particles, in sediments were  $36.237,04 \pm 16.702,59$  kg-1 particles, and in fish were  $275 \pm 95,82$  ind-1 particles or  $654,28 \pm 279,71$  gram-1 particles. The results of the Two Sample T-Test analysis

showed that there was no difference in water, sediment, and mosquitofish (ind-1 particles) between the two situ, but there was a significant difference in mosquitofish (g-1 particles) from the two situ. Fiber and black were respectively the predominant shapes and colors of microplastics in all samples.