A Methodological framework for multi date and multisensor remotelysense image interpretation based on uniform classification scheme = Metodologi interpretasi citra Inderaja multitemporal dan multisensor berdasarkan klasifikasi uniform

Aniati Murni Arymurthy, author

Deskripsi Lengkap: https://lib.ui.ac.id/detail?id=74502&lokasi=lokal

Abstrak

This dissertation describes the synergy use of remote sensing data from multi-temporal and multi sensor (optical and radar) for improving our understanding of the land-cover structural phenomena. A tropical country like Indonesia has a high cloud coverage throughout the year with a maximum during the rainy season, and hence the availability of cloud-free optical images is minimal. To solve this problem, radar images have been intensively introduced. The radar images are cloud-free but their use is hampered due to their speckle noise and topographic distortions, and the lack of a suitable radar image classification system. $\langle br / \rangle \langle br / \rangle$

In many cases, the use of optical or radar image alone is not sufficient. Therefore, the main objectives of this research are: (i) to develop a framework for multi date and multi sensor (optical and radar) image classification; (ii) to solve the cloud cover problem in optical images; and (iii) to obtain a more consistent image classification using multi date and multi sensor images. We have proposed a framework for multi date and multi sensor image classification scheme. The term uniform means that the same procedure can be used to classify optical or radar images, low-level mosaic or fused images, single or multiple feature images.

To be able to conduct a multi temporal and multi sensor analysis, we have unified the optical and radar image classification procedure after finding that both optical and radar images consist of homogeneous and textured regions. A region is considered as homogeneous if the local variance of gray level distribution is relatively low, and a region is considered as textured if the local variance is high. We used a multivariate Gaussian distribution to model the homogeneous part and a multinomial distribution to model the gray level co-occurrences of the textured part, and applied a multiple classifier system to improve the classification accuracy.

The proposed framework incorporates the image mosaicking and data fusion at the low-level stage (before

the classification process) as well as at the high-level stage (after the classification process). For cloud cover removal, the image mosaicking at the low-level stage is usually done using multi temporal optical images, whereas mosaicking at the high-level stage is applied to the classified optical and radar images. To be able to obtain a cloud-free image, we have modified the existing Soofi and Smith algorithm which is using multi temporal optical images to an algorithm using multi sensor images. In the high-level data fusion, we have also been able to incorporate a mechanism for cloud cover removal by omitting the information from the optical sensor and using only the information from the radar sensor. According to a case study in our experiment, the cloud cover removal and image classification using the low-level image mosaicking, the high-level image mosaicking, and the high-level data fusion gave 80.2%, 76.2%, and 80.5% classification accuracy, respectively.

The high-level data fusion combines the decisions from several input images to obtain a consensus of classified image. We have applied both the maximum joint posterior probability and the highest rank method for the decision combination functions. We have utilized two existing data fusion methods and have proposed an alternative data fusion method based on the compound conditional risk. According to the experimental results, the decision combination function based on the maximum joint posterior probability favors the optical feature image, while the highest rank method favors the radar feature image. The preference of using the maximum joint posterior probability results in the domination of optical features in the fusion result, and the classification accuracy of the fused image can be better 8.5% in average than the individual radar classified image.