

Kepadatan dan indeks makanan terbesar kerang kima *tridacna maxima*) di perairan pulau Pasumpahan kota Padang Sumatera Barat = The density and index of preponderance of the giant clam (*tridacna maxima*) in the water Pasumpahan Island, Padang City, West Sumatera

Jabang Nurdin, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=89593&lokasi=lokal>

Abstrak

Telah dilakukan penelitian tentang kepadatan dan indeks makanan terbesar kerang kima *Tridacna maxima* di perairan Pulau Pasumpahan dari bulan Juni-Oktober 2003. Lokasi pencuplikan kerang *T. maxima* dibagi atas tiga strata yaitu strata I (sebelah barat Pulau Pasumpahan yang terdiri dari rata-rata karang dengan padang lamun yang banyak), strata II (sebelah utara Pulau Pasumpahan yang terdiri dari rata-rata karang yang didominasi oleh makroalga) dan strata III (sebelah timur Pulau Pasumpahan yang terdiri dari ekosistem mangrove). Pada masing-masing strata diambil tiga individu kerang *T. maxima*. Kerang yang didapatkan diambil lambungnya kemudian isi lambung kerang *T. maxima* dianalisis. Hasil penelitian didapatkan bahwa komposisi makanan alami kerang *T. maxima* yaitu fitoplankton dan zooplankton. Berdasarkan strata pencuplikan sampel kepadatan makanan alami kerang *T. maxima* terbesar didapatkan pada strata III yaitu 1813,54 individu plankton/individu kerang dan terendah strata II yaitu 1256,52 individu plankton/individu kerang. Kepadatan makanan alami kerang *T. maxima* tertinggi berdasarkan kelas yaitu Bacillariophyceae, kemudian diikuti kelas Crustaceae dan terendah kelas Rotifera, Nilai indeks makanan terbesar yang tertinggi didapatkan dari kelas Bacillariophyceae yaitu 30,47 dan terendah kelas Rotifera yaitu 0,93. Dari nilai indeks makanan terbesar didapatkan bahwa makanan utama kerang *T. maxima* adalah plankton dari kelas Bacillariophyceae dan Crustacea.

The Density And Index Of Preponderance Of The Giant Clam (Tridacna Maxima) In The Water Pasumpahan Island, Padang City, West Sumatera. The research has been done about the density and index of preponderance of the giant clam Tridacna maxima in the water Pasumpahan Island from June to October in 2003. The giant clam T. maxima collection sites in the water Pasumpahan Island were divided three strata i, e. strata I (at west Pasumpahan Island still have coral reef with seagrass), strata II (at north Pasumpahan Island which the coral reef was dominated by macro algae), strata III (at east Pasumpahan Island with mangrove ecosystem). At each strata the giant clam T. maxima were collected three T. maxima. The giant clam was taken its stomach and then the stomach of giant clam were analysed. Results of the research showed that the composition of the natural food giant clam were phytoplankton and zooplankton. Based on the samples collection the highest density from natural food of giant clam T. maxima were found at strata III 1813.54 plankton/shell individu and the lowest density of natural food of giant clam T, maxima were 1256.52 plankton/shell individu at strata II. The highest density of natural food of giant clam T. maxima based on class were Bacillariophyceae and then Crustacea and the lowest density were Rotifera. The highest index of preponderance was found from Bacillariophyceae (30.47) and the lowest were Rotifera (0.93). Food preponderance index was calculated to indicate that the phytoplankton of Bacillariophyceae (30.47) and Crustacea (27.41) were the most preference food by T. maxima.