

## **Latex allergen sensitization reaction in exposed and un-exposed groups as well as associated factors**

R.A.H.I. Ariestina, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=89775&lokasi=lokal>

---

### **Abstrak**

General aim: this study aims to determine the prevalence of sensitization reaction on exposed and unexposed groups of workers at a latex factory. Specific aim: to determine the latex allergen sensitization reaction and factors such as atopy, age, sex, smoking, length of working, exposure concentration.

Sample and method: the sample population consists of workers at a sphymomanometer and condom factory who were, exposed and un-exposed to latex material at the factory. The number of sample was determined based on the two proportion formula.

This study was a cross-sectional study using the Chi square test, Fisher test, Mann Whitney test, and Kruskal Wallis test.

The results of the study and conclusion: The prevalence of latex allergen sensitization reaction in the exposed group was 2.38%, while in the un-exposed group 3.33%. The correlation between exposed and un-exposed group to the latex allergen sensitization reaction (overall) was not significant ( $p=0.181$ ), but if the correlation of each factor was calculated, the most significant correlation was between the negative latex allergen sensitization reaction with a strength of +2 or above ( $p=0.014$ ).

In this study, atopy ( $p=0.000$ ), exposed concentration ( $p=0.014$ ), and smoking ( $p=0.018$ ) are factors that were associated with latex allergen sensitization reaction. Workers at the condom factory were at higher risk than those at the sphymomanometer factory, with a prevalence of 7.14% and 2.31% respectively.