

## Specific subjective symptoms for gastroesophageal reflux disease in ulcer like dyspepsia

Irsan Hasan, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=90096&lokasi=lokal>

---

### Abstrak

**Background:** The determination of branched chain amino acids (BCAA) to tyrosine ratio (BTR) was available in making differentiation of chronic hepatitis from liver cirrhosis, because there was a strong association between BTR and staging (fibrosis) scores. Branched chain amino acids to tyrosine ratio have a correlation with Fischer ratio and the examination is easier because it can be done by enzymatic assay.

**Materials and Methods:** To evaluate the correlation between BTR and Child-Pugh score, we examined the amino gram of 52 liver cirrhosis patients consisted of 26 Child-Pugh A, 19 Child-Pugh B, and 7 Child-Pugh C. The examination of amino gram was done by High Pressure Liquid Chromatograph (HPLC) analyzer. Branched chain amino acids to tyrosine ratio were compared to Child-Pugh score, albumin, ammonia level, number connection test to Fischer ratio.

**Results:** Significant differences in BTR among Child-Pugh A, B, C were observed (Child-Pugh A  $7.75 \pm 1.2$ ; Child Pugh B  $6.0 \pm 1.23$  and Child Pugh C  $4.38 + 3.14$  ( $p = 0.000$ )). Branched chain amino acids to tyrosine ratio had a weak correlation with albumin ( $r = -0.292$ ;  $p = 0.036$ ), ammonia level ( $r = 0.376$ ;  $p = 0.006$ ) and strong correlation with Fischer ratio ( $r = 0.818$ ;  $p = 0.000$ ). There was no significant correlation between BTR and number connection test.

**Conclusion:** These results showed that the determination of the molar ratio of branched chain amino acids to tyrosine well reflected the severity of liver cirrhosis and it can be used as a substitute of Fischer ratio.