

## Nutritional status of under five children and their mothers before and during monetary crisis in east jakarta

Pritasari, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=96584&lokasi=lokal>

---

### Abstrak

#### <b>ABSTRACT</b><br>

Since July 1997, there was a monetary crisis in Indonesia. This condition influences health and nutritional status of the community. There was a decrease of purchasing power, due to lack of job opportunities. This condition led to decreasing in quality and quantity of food intake of children under-five in the community.

<br><br>

Four cross-sectional surveys were carried out in East Jakarta in September 1993, April 1998, December 1998, and March 1999. There were totally 1859 under-five children and 1622 households assessed within the four surveys. Multi-stage random sampling was used in the first survey (September 1993) by Susilowati. Regardless of whether there were under-fives or not, the study covered all sub-districts, selecting households having children under the age of 60 months from 16 village units of 10 sub-districts. And the next three surveys used the same study area as the first study. Data on socio-economic, food consumption, environmental situation, and nutrition status were collected using pre-coded questionnaires and anthropometric measurements. The percentage of fathers without occupation increased in 1999 (7.3%) as compared to 1993 (2%). The anthropometric status of under-five children in September 1993, April 1998, December 1998, and March 1999 shows the prevalence of stunting as follows: 22.7%, 16.1%, 25.4%, and 22% (p value < 0.05, Chi Square). While the prevalence of wasting are as follows: 15%, 17.9%, 11.5%, and 14.1% (p value < 0.05, Chi Square). There were no significant differences in nutritional status between boys and girls within the four surveys. During the monetary crisis, there were significant differences in prevalence of underweight (April '98 survey) in children from lower strata/percentile of per capita food expenditure and upper level of strata/percentile of per capita food expenditure (p < 0.05, Chi Square). Anthropometric indicators were calculated using growth reference data of the National Center for Health Statistics (1977).

<br><br>

The prevalence of stunting and wasting were also significantly higher in the group receiving food aid compared to those without food aid. There is no significant difference in prevalence of malnourished mothers by using cut-off point  $BMJ < 18.5$ . The prevalence of malnourished mothers found in September 1993 was 14.7%, in April 1998 was 13.1%, in December 1998 was 16.2%, and in March 1999 was 13%. The same situation also describes  $BMI > 25$ , which has no changes from 1993 to 1999. The prevalence of overweight ( $BMI > 25$ ) mothers found in September 1993 was 24.2%, April 1998 was 21.3%, December 1998 was 19.1%, and March 1999 was 24.6%. It seems that the crisis has no impact on the nutritional status of non-pregnant mothers.

<br><br>

The crisis was associated with a clear negative shift in household food consumption. The intake of animal protein sources significantly decreased (p value <

0.05.chi-square). as a coping mechanism the plant protein as well as green leafy vegetables consumption increased markedly)

<br><br>

Based on strata/percentile of per capita food expenditure there were no difference within the strata in all surveys in eggs consumption in the family but there were difference for poultry, meat, fish and milk ( $p < 0.05$ . Chi Square)

<br><br>

There was association between children nutritional status and frequency of food intake of animal protein, milk, fruit and cooking oil ( $p$  value  $< 0.05$  anova). There were also association between nutritional status of under live children with food expenditure number of household member, mother nutritional status (BMI and MUAC). Duration of breast-feeding snack consumption, total expenditure and food and program