

## Fasies dan Lingkungan Pengendapan Formasi Cinambo, Daerah Kecamatan Jatinunggal, Sumedang, Jawa Barat = Facies and Depositional Environment of Cinambo Formation, Jatinunggal Area, Sumedang, West Java

Andrea Mutiara Listanti, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=9999920518043&lokasi=lokal>

---

### Abstrak

Lokasi penelitian berada pada Formasi Cinambo wilayah Daerah Jatinunggal, Sumedang Jawa Barat. Pada lokasi ini, tersingkap dengan baik beberapa singkapan sepanjang Anak Sungai Ci Jaweu pada Desa Cibuyung dan Desa Cimanintin yang terbagi menjadi dua lintasan. Litologi penyusunnya adalah feldspathic wacke, allochemic mudstone, foraminiferal-packstone, foraminiferal-wackestone, dan batulanau. Porositas batumannya memiliki rentang poor – excellent (>25%). Berdasarkan pengamatan lapangan, pengukuran stratigrafi, dan analisis petrografi, lokasi penelitian tersusun atas Kelompok Fasies B, C, D, dan G yang terbagi lagi menjadi 7 anggota litofasies. Kelompok-kelompok fasies yang saling berasosiasi kemudian dikelompokkan menjadi 3 asosiasi fasies yaitu Sandy-silt lobe distal, distal levee, dan distal silt-mud lobe. Analisis mikrofosil yang dilakukan, menunjukkan lapisan pada lokasi penelitian memiliki umur Miosen Tengah/Miosen Akhir dan terendapkan pada lingkungan upper-lower bathyal. Formasi Cinambo yang tersingkap pada daerah ini diinterpretasikan berada pada lingkungan kipas laut dalam bagian tengah (middle fan).

.....The research location is in the Cinambo Formation in the Jatinunggal Region, Sumedang, West Java. At this location, several outcrops along the Ci Jaweu River in Cibuyung Village and Cimanintin Village are well exposed and are divided into two tracks. Its constituent lithologies are feldspathic wacke, allochemic mudstone, foraminiferal-packstone, foraminiferal-wackestone, and siltstone. The rock porosity ranges from poor to excellent (>25%). Based on field observations, stratigraphic measurements, and petrographic analysis, the study site is composed of Facies B, C, D, and G which are further divided into 7 lithofacies. Facies groups that are associated with each other are then grouped into 3 facies associations namely distal Sandy-silt lobe, distal levee, and distal silt-mud lobe. The microfossil analysis performed showed that the layers at the study site are Middle Miocene/Lower Miocene and were deposited in an upper-lower bathyal environment. The Cinambo Formation exposed in this area is interpreted to be in the middle fan environment.