

Hubungan Ketahanan Pangan dan Faktor Lainnya dengan Kejadian Underweight pada Anak Usia 24-59 Bulan di Kecamatan Tanjung Priok Jakarta Utara Tahun 2023 = The Relationship between Food Security and Other Factors with the Incidence of Underweight among Children Aged 24-59 Months in Tanjung Priok Subdistrict North Jakarta in 2023

Jihan Farhanah, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=9999920522343&lokasi=lokal>

Abstrak

Latar Belakang: Underweight merupakan masalah gizi yang dapat dialami oleh anak berusia di bawah 5 tahun yang jika tidak segera ditangani dapat mengakibatkan wasting dan stunting (WHO, 2019). Tujuan: Penelitian ini bertujuan mengetahui hubungan ketahanan pangan dan faktor lainnya dengan kejadian underweight pada anak usia 24-59 bulan di Kecamatan Tanjung Priok, Jakarta Utara. Metode: Penelitian ini menggunakan desain studi cross sectional dengan menggunakan pengumpulan data primer pada Juni 2023 dan teknik simple random sampling dengan total sampel sebanyak 184 anak usia 24-59 bulan. Pengumpulan data dilakukan dengan cara pengukuran antropometri, wawancara kuesioner, Semi Quantitative Food Frequency dan food recall 1x24 jam. Analisis data yang dilakukan berupa analisis bivariat menggunakan uji Chi-Square dan multivariat menggunakan uji Regresi Logistik Ganda. Hasil: Hasil penelitian menunjukkan bahwa sebanyak 17,4% anak usia 24-59 bulan di Kecamatan Tanjung Priok mengalami underweight. Berdasarkan analisis bivariat, penelitian ini menunjukkan bahwa terdapat 6 variabel yang berhubungan signifikan dengan underweight pada anak usia 24-59 bulan, yaitu asupan energi ($p\text{-value} = 0,001$), asupan lemak ($p\text{-value} = 0,000$), asupan karbohidrat ($p\text{-value} = 0,011$), riwayat BBLR ($p\text{-value} = 0,010$), ketahanan pangan rumah tangga ($p\text{-value} = 0,023$), dan pengetahuan gizi ibu atau pengasuh ($p\text{-value} = 0,026$). Selain itu, hasil multivariat menunjukkan bahwa riwayat BBLR ($OR=5,1$; $95\%CI=1,564 - 16,866$) sebagai faktor dominan kejadian underweight pada anak usia 24-59 bulan di Kecamatan Tanjung Priok. Kesimpulan: Secara analisis statistik, asupan energi, asupan lemak, asupan karbohidrat, ketahanan pangan rumah tangga dan pengetahuan ibu atau pengasuh berhubungan signifikan dengan kejadian underweight anak usia 24-59 bulan.

.....**Background:** Underweight is a nutritional problem that can be experienced by children under 5 years old. If it persists for a long time and not treated immediately, underweight can result in wasting and stunting (WHO, 2019). **Objective:** This study aims to determine the relationship between food security and other factors with the incidence of underweight in children aged 24-59 months in Tanjung Priok Subdistrict, North Jakarta. **Method:** This research uses a cross-sectional study design with primary data collection in June 2023. The sampling technique used is simple random sampling with a total sample of 184 children aged 24-59 months in Tanjung Priok Subdistrict. Data collection is conducted through anthropometric measurements, questionnaire interviews, Semi Quantitative Food Frequency and 24-hour food recall. The data analysis includes univariate analysis, bivariate analysis using the Chi-Square test, and multivariate analysis using the Multiple Logistic Regression test. **Results:** This research shows that 17,4% of children aged 24-59 months in Tanjung Priok Subdistrict experience underweight. Based on bivariate analysis, this study indicates that there are six variables significantly associated with underweight in children aged 24-59 months, namely energy intake ($p\text{-value} = 0,001$), fat intake ($p\text{-value} = 0,000$), carbohydrate intake ($p\text{-value} =$

0,011), history of low birth weight (p-value = 0,010), household food security (p-value = 0,023), and maternal or caregiver nutrition knowledge (p-value = 0,026). Furthermore, the multivariate results show that a history of low birth weight (OR=5,1; 95%CI=1,564 – 16,866) is the dominant factor for underweight in children aged 24-59 months in Tanjung Priok Subdistrict. Conclusion: Based on statistical analysis, energy intake, fat intake, carbohydrate intake, household food security, and maternal or caregiver nutrition knowledge are significantly associated with underweight in children aged 24-59 months.