

Proporsi gambaran penyakit paru interstisial pada pasien kanker paru karsinoma bukan sel kecil dengan terapi epidermal growth factor receptor tyrosine kinase inhibitor di RSUP Persahabatan = The proportion of interstitial lung disease features in patients with non small cell lung cancer treated with epidermal growth factor receptors tyrosine kinase inhibitor at Persahabatan Hospital

Marscha Iradyta Ais, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=9999920522800&lokasi=lokal>

Abstrak

Latar Belakang: Jumlah kasus KPKBSK diperkirakan 85% dari seluruh kasus kanker paru dan 40% diantaranya adalah jenis adenokarsinoma. Sebanyak 10%-30% pasien adenokarsinoma mengalami mutasi EGFR dan mendapatkan terapi EGFR-TKI. Mayoritas pasien KPKBSK memiliki respons dan toleransi baik terhadap terapi EGFR- TKI tetapi sebagian kecil pasien mengalami penyakit paru interstisial akibat EGFR-TKI. Penelitian ini bertujuan untuk mengetahui proporsi gambaran penyakit paru interstisial pada pasien KPKBSK dengan terapi EGFR-TKI di RSUP Persahabatan.

Metode: Penelitian ini merupakan penelitian observasional analitik dengan pendekatan kohort retrospektif yang dilakukan bulan Januari 2021 hingga Juni 2022. Subjek penelitian adalah pasien KPKBSK yang mendapatkan terapi EGFR-TKI. Subjek penelitian dipilih sesuai kriteria inklusi dan eksklusi. Pengambilan data melalui data sekunder berupa rekam medis dan hasil CT scan toraks pasien yang kontrol di poliklinik onkologi RSUP Persahabatan.

Hasil: Pada penelitian ini diperoleh 73 subjek penelitian, pasien KPKBSK dengan mutasi EGFR yang mendapatkan terapi EGFR-TKI di RSUP Persahabatan. Sebanyak 12 dari 73 subjek penelitian mengalami gambaran ILD yang dievaluasi berdasarkan CT scan toraks RECIST I dan II dengan karakteristik jenis kelamin laki-laki (22,2%), kelompok usia 40-59 tahun (19,4%), perokok (24,1%), indeks brinkman berat (42,9%) dan mendapatkan terapi afatinib (26,1%). Proporsi gambaran ILD pada pasien KPKBSK dengan terapi EGFR-TKI adalah opasitas retikular (58,3%), parenchymal band (33,3%), ground-glass opacities (25%), traction bronchiectasis (25%) dan crazy paving pattern (8,3%). Hasil analisis bivariat dan multivariat menunjukkan tidak terdapat perbedaan antara faktor-faktor seperti jenis kelamin, usia, jenis EGFR-TKI, riwayat merokok, indeks brinkman, riwayat penyakit paru dan tampilan status terhadap gambaran ILD.

Kesimpulan: Gambaran ILD pada pasien KPKBSK dengan terapi EGFR-TKI meliputi opasitas retikular, parenchymal band, ground-glass opacities, traction bronchiectasis dan crazy paving pattern. Tidak terdapat perbedaan bermakna secara statistik antara faktor-faktor yang memengaruhi terhadap gambaran ILD.

.....**Background:** The number of cases of NSCLC is estimated around 85% of all lung cancer cases and 40% among them are adenocarcinoma. Approximately 10%-30% of adenocarcinoma patients have EGFR mutations and receive EGFR-TKI therapy. The majority of NSCLC patients have a good response and tolerance to EGFR-TKI therapy, but a small group of patients experience EGFR-TKI induced interstitial lung disease. This study aims to determine the proportion of features of interstitial lung disease in NSCLC patients treated with EGFR-TKI at Persahabatan Hospital.

Methods: This study was an analytic observational with a retrospective cohort approach that was conducted from January 2021 until June 2022. The subject were NSCLC patients who received EGFR-TKI treatment.

The inclusion and exclusion criteria were used to determine which subjects will be included in the study. Data collection through secondary data from medical record and chest CT scan results of patients controlled at oncology polyclinic at Persahabatan Hospital.

Result : In this study, there were 73 subjects of NSCLC with EGFR mutations and received EGFR-TKI therapy at Persahabatan Hospital. There were 12 out of 73 subjects had ILD features which were evaluated based on RECIST I and II chest CT scan with predominant of male (22.2%), age group 40-59 years old (19.4%), smokers (24.1%), severe Brinkman index (42.9%) and received afatinib (26.1%). The proportion of ILD features in NSCLC patients with EGFR-TKI therapy are reticular opacities (58.3%), parenchymal bands (33.3%), ground-glass opacities (25%), traction bronchiectasis (25%) and crazy paving pattern (8.3%). The results of bivariate and multivariate analyzes showed that there was no differences between factors such as sex, age, type of GEFR-TKI, smoking history, Brinkman index, history of lung disease and performance status with features of ILD.

Conclusion: Features of ILD in NSCLC patients with EGFR-TKI therapy include reticular opacities, parenchymal bands, ground-glass opacities, traction bronchiectasis and crazy paving pattern. There is no statistically significa