

Faktor - Faktor yang Berhubungan dengan Kematian Akibat Difteri di Indonesia Tahun 2018 = Factors Associated with Diphtheria Death in Indonesia 2018

Nurhanifah Hamdah, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=9999920532987&lokasi=lokal>

Abstrak

Difteri merupakan penyakit re-emerging yang disebabkan oleh bakteri *Corynebacterium diphtheria* dan menyebabkan kematian. Kematian akibat difteri ini di dunia memiliki CFR 5 – 10% dan di Indonesia CFR difteri sebesar 2% sehingga masih menjadi masalah kesehatan masyarakat. Penelitian ini bertujuan untuk mengetahui faktor-faktor yang berhubungan dengan kematian akibat difteri di Indonesia Tahun 2018. Penelitian ini menggunakan data surveilans laporan rutin kasus difteri di Indonesia tahun 2018. Desain studi yang digunakan adalah cross-sectional dengan unit analisis individu yang menderita difteri di Indonesia tahun 2018. Pada penelitian ini didapatkan 817 kasus dengan 2,8% penderita meninggal dan tempat tinggal responden paling banyak di Jawa Barat, yaitu 211 penderita. Faktor jenis kelamin, status imunisasi primer difteri, dan status pemberian Anti Difteri Serum (ADS) tidak berhubungan secara statistik dengan kematian akibat difteri. Faktor yang berhubungan secara statistik dengan kematian akibat difteri, yaitu umur < 15 tahun (OR = 7,863; 95% CI = 1,831 – 33,77) dan diagnosis penderita difteri konfirmasi laboratorium (OR = 2,774; 95% CI = 1,000 – 7,693).

.....Diphtheria is a re-emerging disease caused by the *Corynebacterium diphtheria* bacteria and causes death. The death from diphtheria in the world has CFR of 5-10% while in Indonesia has CFR is 2% that make it becomes one of public health concerns. This study aims to determine the factors associated with diphtheria deaths in Indonesia in 2018 using surveillance data on routine reports of diphtheria cases in Indonesia in 2018. The study design used was cross-sectional with a unit of analysis of individuals suffering from diphtheria in Indonesia in 2018. In this study there were 817 cases with 2.8% of patients dying and mostly of them living in West Java province (211 patients). Sex factors, primary diphtheria immunization status, and Serum Anti Diphtheria (ADS) status were not statistically related to diphtheria deaths. Factors statistically associated with diphtheria deaths were age <15 years (OR = 7.863; 95% CI = 1,831 - 33,77) and diagnosis of diphtheria patients from laboratory confirmation (OR = 2,774; 95% CI = 1,000 - 7,693).