

Korelasi Kerapatan Mangrove dengan Kelimpahan Makroplastik di Suaka Margasatwa Pulau Rambut, DKI Jakarta = Correlation of Mangrove Density with Macroplastic Abundance in Pulau Rambut Wildlife Reserve, DKI Jakarta

Hana Restu Oktavia, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=9999920534493&lokasi=lokal>

Abstrak

Produksi plastik yang terus meningkat disertai sistem pengelolaan yang buruk menyebabkan tercemarnya lingkungan oleh limbah plastik. Pencemaran tersebut juga terjadi pada lautan dan daerah pesisir. Vegetasi mangrove di daerah pesisir dapat berperan sebagai perangkap bagi limbah plastik. Plastik-plastik yang terperangkap biasanya merupakan plastik dengan ukuran besar atau biasa disebut dengan makroplastik. Penelitian ini dilakukan untuk mengetahui korelasi antara kerapatan mangrove jenis pohon, pancang, dan semai dengan kelimpahan makroplastik di Suaka Margasatwa Pulau Rambut, DKI Jakarta. Pengambilan data kerapatan mangrove dilakukan pada plot kuadran 10 x 10 m yang tersebar di 30 titik pada 7 stasiun pengamatan. Pengambilan data kelimpahan makroplastik dilakukan pada sub-plot kuadran berukuran 1 x 1 m yang berada di dalam plot kuadran mangrove dengan 3 kali pengulangan. Data-data tersebut kemudian dianalisis menggunakan uji korelasi Spearman. Terdapat korelasi signifikan yang bersifat positif sangat kuat antara kerapatan mangrove jenis pohon dengan kelimpahan makroplastik di Suaka Margasatwa Pulau Rambut. Kerapatan mangrove jenis pancang dengan kelimpahan makroplastik di Suaka Margasatwa Pulau Rambut memiliki korelasi yang tidak signifikan dan bersifat negatif sangat lemah. Terdapat korelasi signifikan yang bersifat positif lemah antara kerapatan mangrove jenis semai dengan kelimpahan makroplastik di Suaka Margasatwa Pulau Rambut.

.....Plastic production continues to increase accompanied by poor management systems that causes environmental pollution by plastic waste. Pollution also occurs in the oceans and coastal areas. Mangrove vegetation in coastal areas can act as a trap for plastic waste. Trapped plastics are usually large plastics or commonly known as macroplastics. This research was conducted to determine the correlation between the density of mangrove tree species, saplings, and seedlings with macroplastic abundance in Pulau Rambut Wildlife Reserve, DKI Jakarta. Mangrove density data was collected in 10 x 10 m quadrant which was spread over 30 points at 7 observation stations. Data collection on macroplastics abundance was carried out in the 1 x 1 m sub-plot quadrant which placed inside the mangrove quadrant plot with 3 repetitions. These data were then analyzed using the Spearman correlation test. There is a significant positive and very strong correlation between the density of mangrove tree species with the macroplastics abundance in Pulau Rambut. The density of sapling mangroves with macroplastic abundance in Pulau Rambut has an insignificant correlation and is very weakly negative. There is a significant correlation that is weak positive between the density of mangrove seedlings and the macroplastics abundance in Pulau Rambut.