

Perbandingan pemberian ketamin dan lidokain untuk mengurangi derajat nyeri pada saat induksi anestesi menggunakan propofol di RS Saiful Anwar Malang = Comparison of ketamine and lidocaine to reducing pain during anesthesia induction with propofol in Saiful Anwar Hospital Malang

Nopian Hidayat, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=9999920535557&lokasi=lokal>

Abstrak

Latar Belakang. Propofol merupakan obat anestesi intravena yang paling sering digunakan dalam pembiusan umum tetapi propofol dapat menimbulkan rasa nyeri pada lokasi injeksi dengan angka kejadian 28-90%. Pemberian lidokain sebelumnya paling sering digunakan untuk mengurangi nyeri yang ditimbulkan propofol, akan tetapi tingkat kegagalannya 13-32. Tujuan penelitian ini adalah untuk mengetahui perbandingan pemberian pre-emptive ketamin 0,1 mg/kg dan lidokain 1 mg/kg untuk mengurangi derajat nyeri pada saat induksi anestesi menggunakan propofol.

Metode. Penelitian ini merupakan uji klinis tersamar ganda, bersifat eksperimental. Pasien dengan kriteria klinis ASA I-II sejumlah 50 orang yang akan menjalani operasi elektif dengan pembiusan umum, dilakukan randomisasi sederhana menjadi 2 kelompok yaitu kelompok I (lidokain 1 mg/kg) dan kelompok II (ketamin 0,1 mg/kg) yang diberikan 1 menit sebelum induksi propofol. Derajat nyeri dinilai berdasarkan Verbal Rating Scale (VRS).

Hasil. Penelitian menunjukkan pemberian pre-emptive ketamin dapat menurunkan derajat nyeri yang lebih baik (84% tidak nyeri, 16% nyeri ringan) dibandingkan kelompok pre-emptive lidokain (56% tidak nyeri, 28% nyeri ringan, 12% nyeri sedang dan 4% nyeri berat) dengan nilai $p = 0.021$ (p bermakna < 0.05) pada uji statistik menggunakan Mann Whitney.

Kesimpulan. Pemberian pre-emptive ketamin 0,1 mg/kg BB intravena lebih baik dibandingkan dengan pemberian pre-emptive lidokain 1 mg/kg BB untuk mengurangi derajat nyeri akibat penyuntikan propofol intravena.

.....

Background. Propofol is a popular IV anesthetic induction drug that causes pain when given IV. The incidence of which is between 28-90%. Lidocaine pre-treatment has been commonly proposed to decrease propofol induced pain, but its failure rate is between 13-32%. The purpose of this study was to compare a pre-emptive ketamine 0,1 mg/kg and pre-emptive lidocaine 1 mg/kg to minimize the injection pain of propofol during anesthesia induction.

Methods. A comparative, randomized, double blind study of 50 patients (ASA I-II) scheduled surgery under general anesthesia were randomly allocated into two groups. Group I received lidocaine 1 mg/kg and group II received ketamine 0,1 mg/kg one minute before the anesthesia induction with propofol IV. Each patient's pain score were evaluated by using Verbal Rating Scale (VRS)

Result. The result of this study described that pre-emptive ketamine had significantly lower incidence of pain and lower pain score (84% no pain, 16% mild pain) compared with pre-emptive lidocaine (56% no pain, 28% mild pain, 12% moderate pain and 4% severe pain) with p value = 0.021 (significant $p < 0.05$) using Mann Whitney statistic test.

Conclusion. Pre-emptive ketamine 0,1 mg/kg significantly in reducing degree of propofol pain injection compare with pre-emptive lidocaine 1 mg/kg IV.