

Analisis Asuhan Keperawatan Keperawatan Pada Pasien Efusi Pleura Bilateral dengan Penerapan Intervensi Posisi High Fowler dan Orthopneic Sebagai Upaya Memperbaiki Pola Napas = Analysis of Nursing Care for Bilateral Pleural Effusion Patients with the Implementation of High Fowler and Orthopneic Position Interventions as an Effort to Improve Breathing Patterns

Dea Putri Amalia, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=9999920539053&lokasi=lokal>

Abstrak

Efusi pleura merupakan kejadian akumulasi cairan yang berada diantara lapisan parietal dan viseral. Penumpukan cairan pada rongga pleura akan menyebabkan masalah utama pada sistem pernapasan. Studi ini bertujuan untuk menganalisis intervensi keperawatan mandiri berupa pengaturan posisi high fowler dan orthopneic pada pasien dengan masalah pola nafas tidak efektif pada kasus efusi pleura. Metode yang digunakan adalah studi literatur terkait pemberian posisi high fowler dan orthopneic pada pasien yang memiliki masalah pernapasan. Hasil studi menunjukan pemberian posisi high fowler dan orthopneic keduanya berdampak pada peningkatan saturasi oksigen dan penurunan frekuensi napas pada kondisi sesak. Analisis studi ini merekomendasikan pengaturan posisi high fowler dan orthopneic efektif untuk diimplementasikan pada pasien dengan masalah pola napas tidak efektif.

..... Pleural Effusion is an accumulation of fluid between the parietal and visceral layers. the buildup of fluid in the pleural cavity will cause major problems with the respiratory system. this study aims to analyze independent nursing interventions in the form of setting high fowler and orthopneic positions in patients with problems with ineffective breathing patterns in cases of pleural effusion. the method used is a literature study regarding the provision of high fowler and orthopneic positions in patients who have breathing problems. the results of the study show that providing the high fowler and orthopneic positions has an impact on increasing oxygen saturation and reducing respiratory frequency in shortness of breath. the analysis of this study recommends that high fowler's and orthopneic positions be effective to be implemented in patients with ineffective breathing pattern problems.