

Model Geographically Weighted Regression Determinan Angka Kematian Neonatal Di Provinsi Jawa Tengah, Jawa Timur, Jawa Barat Dan Daerah Istimewa Yogyakarta Tahun 2022 = Geographically Weighted Regression Model Determinants of Neonatal Mortality Rates in the Provinces of Central Java, East Java, West Java and the Special Region of Yogyakarta in 2022

Mochamad Anwarid Ardans Pratama, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=9999920548536&lokasi=lokal>

Abstrak

Bayi usia dibawah satu bulan termasuk kedalam kategori usia dengan risiko gangguan kesehatan paling tinggi, risiko tersebut dapat berakibat fatal yaitu kejadian kematian apabila tidak mendapat pelayanan kesehatan yang memadai. Kematian neonatal di Provinsi Jawa Tengah, Provinsi Jawa Timur dan Provinsi Jawa Barat tersebut pada tahun 2022 mencapai 38,38% dari jumlah kematian neonatal pada tingkat nasional. Di Provinsi Daerah Istimewa Yogyakarta selama 4 tahun cenderung meningkat, dengan peningkatan terbesar terjadi di Tahun 2022 dengan peningkatan sebesar 16% dibandingkan Tahun 2021. Tujuan dari penelitian ini adalah untuk mengetahui determinan AKN yang dipengaruhi oleh kondisi spasial di Provinsi Jawa Tengah, Jawa Timur, Jawa Barat dan Daerah Istimewa Yogyakarta pada tahun 2022. Penelitian berdesain studi ekologi dengan unit analisis 105 kabupaten/kota di Provinsi Jawa Tengah, Jawa Timur, Jawa Barat dan Daerah Istimewa Yogyakarta. Data yang dianalisis merupakan data agregat yang bersumber dari data sekunder berupa Profil Kesehatan Provinsi Tahun 2022 dan hasil Susenas Tahun 2022 yang dipublikasikan oleh BPS. Pemodelan dengan metode GWR diketahui bahwa dari 6 variabel yang dianalisis dalam model, terdapat 3 variabel yang berpengaruh signifikan secara statistik dengan AKN yaitu variabel BBLR, kepadatan jalan, dan KN lengkap.

.....Babies aged under one month are included in the age category with the highest risk of health problems, this risk can have fatal consequences, namely death if they do not receive adequate health services. Neonatal deaths in Central Java Province, East Java Province and West Java Province in 2022 will reach 38.38% of the number of neonatal deaths at the national level. In the Yogyakarta Special Region Province, it has tended to increase over the past 4 years, with the largest increase occurring in 2022 with an increase of 16% compared to 2021. The aim of this research is to determine the determinants of AKN which are influenced by spatial conditions in the Provinces of Central Java, East Java, Java West and Special Region of Yogyakarta in 2022. The research has an ecological study design with analysis units of 105 districts/cities in the Provinces of Central Java, East Java, West Java and the Special Region of Yogyakarta. The data analyzed is aggregate data sourced from secondary data in the form of the 2022 Provincial Health Profile and the results of the 2022 Susenas published by BPS. Modeling using the GWR method shows that of the 6 variables analyzed in the model, there are 3 variables which have a statistically significant effect on AKN are the variables LBW, road density, and complete neonatal visit.