

Hubungan Durasi Fototerapi dengan Nilai Bilirubin Total pada Pasien Ikterik Neonatorum di RS Polri Tipe C DKI Jakarta = Correlation of Phototherapy Duration with Total Bilirubin Values in Neonatal Jaundice Patients at Type C Police Hospital DKI Jakarta

Annisa Istiqomah, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=9999920566472&lokasi=lokal>

Abstrak

Ikterus neonatorum merupakan kondisi pada neonatus karena peningkatan bilirubin dalam darah. Ikterus neonatus mempunyai tanda diantaranya kulit dan sklera yang kuning akibat bilirubin tak terkonjugasi. Salah satu terapi hyperbilirubinemia adalah fototerapi. Tujuan penelitian adalah untuk mengetahui hubungan durasi fototerapi dengan nilai bilirubin total akhir pada pasien ikterik neonatorum di RS Polri DKI Jakarta. Penelitian ini menggunakan desain cross sectional dengan total 103 data yang mengalami ikterus neonatorum sebagai responden. Hasil penelitian menunjukkan rata-rata kadar bilirubin awal 14, 81 mg/dL, kadar bilirubin setelah fototerapi selama 24 jam menghasilkan penurunan sebesar 2,60 mg/dL dan fototerapi selama 48 jam menghasilkan rata-rata penurunan bilirubin sebesar 3,69 mg/dL. Terdapat hubungan durasi fototerapi dengan nilai bilirubin total pada 24 jam dan 48 jam pasca fototerapi (p Value $<0,005$).

Rekomendasi penelitian ini adalah perlu dilakukannya eksplorasi faktor-faktor yang berhubungan dengan efektivitas fototerapi.

.....Neonatal jaundice is a condition in neonates due to increased bilirubin in the blood. Neonatal jaundice has signs including yellow skin and sclera due to unconjugated bilirubin. One of the therapies for hyperbilirubinemia is phototherapy. The purpose of the study was to determine the relationship between the duration of phototherapy and the final total bilirubin value in neonatal jaundice patients at DKI Jakarta Police Hospital. This study used a cross sectional design with a total of 103 data who experienced neonatal jaundice as respondents. The results showed an average initial bilirubin level of 14, 81 mg/dL, bilirubin levels after phototherapy for 24 hours resulted in a decrease of 2.60 mg/dL and phototherapy for 48 hours resulted in an average decrease in bilirubin of 3.69 mg/dL. There is a relationship between the duration of phototherapies with total bilirubin values at 24 hours and 48 hours after phototherapy (p value <0.005). The recommendation of this study is the need to explore the factors associated with the effectiveness of phototherapy.