

Effects of Chlorophyll in Papaya Leaves on Superoxide Dismutation and Blood Glucose Level of Diabetic Rats

Choirun Nissa^{1*}, Martha Irene Kartasurya², Banundari Rahmawati³

1. Department of Clinical Pathology, Faculty of Medicine, Universitas Diponegoro, Semarang 50277, Indonesia

2. Department of Nutrition, Faculty of Medicine, Universitas Diponegoro, Semarang 50277, Indonesia

3. Department of Public Health Nutrition, Faculty of Public Health, Universitas Diponegoro, Semarang 50277, Indonesia

*E-mail: ni55a_gizi@yahoo.com

Abstract

Hyperglycemia in diabetes mellitus results in oxidative stress and increases complication development. Experimental studies have shown that chlorophyll has antioxidant activity and papaya leaves contained chlorophyll more than the other green vegetables. This study aimed to evaluate the antioxidant and hypoglycemic role in chlorophyll rich in papaya leaves on diabetic rats. Thirty six rats were randomly divided into 4 groups: without treatment (technique control/TC), diabetic (negative control/NC), diabetic with treatment A (100.3 mg/200 g BW of extract) and diabetic with treatment B (200.6 mg/200 g BW of extract). Diabetic induction was conducted by injecting streptozotocin 40 mg/kg BW intraperitoneally. Extract was given by nasogastric tube. Blood glucose level was measured using enzymatic colorimetric GOD-PAP test at before, after 20 and 40 days of treatment. Hepatic superoxide dismutation (SOD) level was measured after 40 days of treatment. Blood glucose levels in 3 diabetic groups were significantly raised after seven days of induction. In Anova and post hoc LSD analysis, both treatments had lower hepatic SOD level than TC ($p = 0.01$), and blood glucose level also decreased after given the treatment ($p = 0.01$). Treatment B had a better antioxidant and hypoglycemic role than treatment A.

Abstrak

Efek dari Klorofil di Daun Pepaya pada Dismutasi Superoksida dan Kadar Glukosa Darah Tikus Diabetes. Hiperglikemia pada pasien diabetes melitus menyebabkan stres oksidatif dan meningkatkan pengembangan komplikasi. Berbagai penelitian eksperimental telah memperlihatkan bahwa klorofil memiliki aktivitas antioksidan dan daun pepaya mengandung klorofil lebih banyak dari sayuran hijau lainnya. Penelitian ini bertujuan untuk mengevaluasi peran antioksidan dan hipoglikemik pada kandungan klorofil di daun pepaya pada tikus diabetes. Tiga puluh enam tikus secara acak dibagi menjadi 4 kelompok: tanpa pengobatan (kontrol teknik/TC), diabetes (kontrol negatif/NC), diabetes dengan pengobatan A (100,3 mg /200 g BB ekstrak), dan diabetes dengan pengobatan B (200,6 mg/200 g BW ekstrak). Induksi diabetes dilakukan dengan menyuntikkan streptozotocin 40 mg/kg BW secara intraperitoneal. Ekstrak diberikan melalui tabung nasogastrik. Kadar glukosa darah diukur dengan menggunakan uji enzim kolorimetri GOD-PAP di sebelum, setelah 20 dan 40 hari pengobatan. Tingkat dismutasi superoksida (SOD) hati diukur setelah 40 hari pengobatan. Kadar glukosa darah pada 3 kelompok diabetes meningkat secara signifikan setelah tujuh hari induksi. Dalam analisa Anova dan uji lanjut LSD, kedua kelompok dengan pengobatan memiliki tingkat SOD hati lebih rendah dari TC ($p = 0,01$) dan kadar glukosa darah juga menurun setelah diberi pengobatan ($p = 0,01$). Pengobatan B memiliki peran antioksidan dan hipoglikemik lebih baik daripada pengobatan A.

Keywords: blood glucose, chlorophyll, diabetes mellitus, papaya leaves, superoxide dismutation

Introduction

Diabetes Mellitus is a non-communicable disease that is closely related to a variety of oxidative activities. It leads to an autoxidation of glucose, glycosylated

proteins, and activation of polyol metabolic pathway that accelerates the formation of reactive oxygen compounds. Furthermore, the formation of reactive oxygen compounds reduces the antioxidant enzyme superoxide dismutase including catalase, ceruloplasmin,

and glutathione which initiated the formation of oxidative stress by increasing the risk of complications.¹ In experimental animal studies, the state of diabetes can be performed using streptozotocin-induced rats.

Increased consumption of phytochemicals that have antioxidant activity can suppress the development of diabetes. In contrast to other phytochemical compounds such as phenolic components, chlorophyll is present in large quantities in plants (\pm 1% dry weight); therefore, it is potential to be one of functional food.² The role of chlorophyll as hypoglycemic agent occurs through inhibition of free radical mechanism. Chlorophyll is an antioxidant chain breaker which donates its electrons to free radicals and forms complexes with peroxy radicals and generated stable product.³ Besides chlorophyll, chloroplasts also contain exogenous superoxide dismutase⁴ and ascorbic acid⁵ which also have antioxidant capacity.

Nowadays, there are a lot of imported food supplements based on chlorophyll. Moreover, papaya leaves which has the highest chlorophyll content among green vegetables, are also very common in Indonesia.⁶ The leaves also contain saponins, vitamins, minerals.⁷ A research showed papaya leaves increase hypoglycemic effect of rats treated with metformin.⁸ The purpose of this study was to analyze the effect of chlorophyll in carica papaya Linnaeus leaves on superoxide dismutase and fasting blood glucose levels of diabetic rats.

Methods

The extraction procedure of chlorophyll in papaya leaves was based on research conducted by Setiary and Nurchayati (2009) using 85% acetone solution with a ratio of the sample and acetone 1:100. The extract was filtered by filter paper and analyzed using UV-Vis spectrophotometer at a wavelength of 644 nm and 663 nm. Setiary and Nurchayati found that the chlorophyll content in papaya leaves is 29.5975 mg/g.

Recommendation of 120 g/day of fruit and vegetable consumption (WNPG) resulted in 150 g wet papaya leaves after conversion with Food Processor II and equivalent to 78 g dry leaves. After being extracted with acetone, the weight was reduced to 10% or 7.8 grams. The calculation of the dose conversion factors for human was based on Gosh.⁹ A safe dose for human weighed 50 kg is equivalent with $(7.8 \times 0.018 \times 50 / 70) / 200$ g of rat body weight, or 100.3 mg/200 g body weight for treatment A, and twofold higher for treatment B. The extract was given 1 ml orally.

A randomized controlled trial over 41 days was conducted at PAU Center for the Study of Food and Nutrition, Gadjah Mada University. After 7 days of adaptation period fed with Comfeed AD II and provided

with drink *ad libitum*, a total of 36 Sprague Dawley rats were randomly divided into 4 groups: 1 group as control with no treatment, other groups are negative control (diabetic with no treatment), treatment A, and treatment B (twofold higher). Streptozotocin was administered intraperitoneally with a dose of 40 mg/kg body weight, which were given to 3 groups. Rats were categorized as having diabetes mellitus when the blood glucose levels were higher than 200 mg/dL.¹⁰

The examination of blood glucose levels was conducted quantitatively for 3 times: pretest (7 days after induction), post-test I (day 21 of treatment), and post test II (day 41 of treatment) with Enzymatic Colorimetric Test method "GOD - PAP". Blood was taken via the orbital sinus. After the second post test of blood glucose level, all rats were terminated using intramuscular ketamine hydrochloride with a dose of 60 mg/kg body weight.¹¹ The measurement of superoxide dismutation levels from rat liver was based on Marklund and Marklund's research,¹² which found that superoxide dismutation enzyme inhibits autooxidation process of pyrogallol into purpurogalin by capturing the oxygen.

Analyses of variance (Anova) was used to assess the difference in superoxide dismutation level of rat liver and the changes of pre and post blood sugar level among 3 groups (negative control, treatment A and treatment B). If the differences were present, Post Hoc LSD were used to assess which groups were different. Data were presented as means \pm SD with all statistical comparisons made at the 0.05 level of significance (Table 1).

This research passed the study of health research ethics committee, Faculty of Medicine, University of Diponegoro, with ethical clearance number 220/EC/FK/RSDK/2012 dated June 18th 2012.

Results and Discussion

Weight, blood glucose and liver weight. The mean weight at baseline, before induction, and before treatments showed no difference. The streptozotocin induction in 3 groups resulted in some unique characteristics of diabetes such as not physically active, polyuria, weight loss, but without polyphagy. The intakes were relatively the same for all groups (isocaloric) at 17.6 gram/day for 40 days, equivalent to 286 kcal/day. After 40 days of treatments, the best growth and development occurred in treatment B (0.93 g/day), while the biggest weight loss presented in negative control group (-0.36 g/day or 6.8%). Without induction (control techniques), the weight gain could reach 1.27 g/day.

Blood glucose levels increased after 7 days of induction and exceeded Braunwald's (2008) criteria (higher than

200 mg/dL). The treatments (21 and 40 days) resulted in lower blood glucose level; even treatment B reached the normal range of 50-125 mg/dL¹³ on day 41, although not as good as normal rat without induction (Figure 1).

The induction also raised the liver weight by 61% (negative control), 44% (treatment A), and 14% (treatment B) compared to control technique. Group treated with a dose of 200.6 mg/200 g body weight have liver weight closer to normal (control technique).

Effect of chlorophyll extract on superoxide dismutation level and blood glucose level. Extract of chlorophyll from papaya leaves was considered to be safe, even with twofold higher dose (treatment B). It tended to have better antioxidant activity, but it still required further histopathologic analysis and toxicity test. Until now, experts have not provided recommended daily dose of chlorophyll.¹⁴ Figure 3 shows that the Negative control group (no treatment) had the highest level of superoxide dismutase (Table 2). The best effect on blood glucose level occurred at day 41th in Treatment B group. Treatment A decreased blood glucose level by 89.83 mg/dL while treatment B by 110.94 mg/dL after 40 days (Figure 2).

Weight, blood glucose and liver weight. There are several implications generated by the induction of streptozotocin. The effects of streptozotocin is irreversible, and causes degranulation and capacity lost in insulin production that leads to hyperglycemia.¹⁵

Streptozotocin also impairs glucose oxidation as well as causing the loss of adipocytes¹⁶ and muscle atrophy accompanied by a decrease in skeletal muscle mass and a loss of structural proteins resulting in weight loss.¹⁷ Structural proteins are proteins that play a role in structural biology support or build living things.¹⁸ Several factors determine the severity of experimental diabetogenic activity, which are the dosage and administration of streptozotocin, animal species, strain, and age of experimental animals.¹⁹

Other effects are not only limited to the increase in blood glucose levels and weight loss, but also an increase in liver weight. Liver is the main organ in detoxification and oxidative processes. Even at the early stages of the disease, a marker of oxidative stress in the liver began to increase. Free radicals are the results of streptozotocin, liver damage, as well as pancreatic β cells.²⁰

Negative control group had the highest level of superoxide dismutation. Higher superoxide dismutation activity means more radicals to neutralized^{2,21} and greater volume of superoxide dismutation to inhibit the autooxidation process of pyrogallol compounds into purpurogalin by capturing oxygen.²² Superoxide dismutase is the highest antioxidant enzymes in the body and are mostly located in liver.²³ This enzyme catalyzes radical superoxide (O₂)²⁴ which in this case comes from streptozotocin.

Table 1. Means of Weight, Blood Glucose Level, and Liver Weight during Study

Indicator	Technique control	Groups		
		Negative control	Treatment A	Treatment B
Weight (g)				
Initial adaptation	206.2±9.74	218.7±10.72	195.4±13.19	199.6±7.43
After induction	213.6±9.64	225.8±10.59	202.6±13.32	205.9±7.66
Initial treatment	224.2±9.54	221.4±10.65	197.9±13.19	201.7±7.73
End of treatment	274.9±9.50	206.4±10.36	221.9±12.69	239.0±7.63
Changes during treatment	50.7±0.47	-15.0±0.78	24.0±0.60	37.3±0.65
Blood Glucose level (mg/dl)				
Baseline (Day 1)	73.0±0.45	223.7±1.16	223.1±0.93	227.2±1.66
Middle (Day 21 st)	74.7±0.60	226.2±1.28	170.7±1.03	140.1±1.07
End (Day 41 st)	75.7±0.65	228.2±1.10	133.3±2.58	116.3±3.30
Liver weight (g/100 gr BB)	3.6±0.17	5.8±0.44	5.2±0.34	4.1±0.13

Table 2. Effect of Chlorophyll in Papaya Leaves on Superoxide Dismutation and Blood Glucose Level

Indicator	Technique control	Groups			p
		Negative control	Treatment A	Treatment B	
Liver superoxide Dismutation level (%)	23.6±0.84 ^a	57.2±1.17 ^b	42.9±0.59 ^c	33.5±0.42 ^d	0.01*
Changes of blood glucose level (mg/dl)	2.7±0.38 ^a	4.5±0.45 ^b	-89.8±2.83 ^c	-110.9±3.45 ^d	0.01*

p = Anova in 3 induction groups (negative control, Treatment A and B)

* = Anova, significant

The average value followed by different letters in the same row indicates significant differences (p <0.05)




Figure 1. Graphic of Rats Blood Glucose Level




Figure 2. Graphic of Blood Glucose Changing




Figure 3. The Effect of Papaya Leaf Chlorophyll Extract on Superoxide Dismutation Level

Higher dose of chlorophyll extract had better antioxidant activity than the lower dose, as indicated from the percentage of inhibition levels were considered to approach the value of control technique. It might be caused by 3 factors. Firstly, the antioxidant properties of chlorophyll. Chlorophyll is a chain breaker antioxidant that works by donating electron to free radicals, especially at the initiating stage before the hydroperoxide is formed. The essential structure of the chlorophyll, which has antioxidant activity, is in the porphyrin structure. This role is supported by magnesium at the core of chlorophyll, primarily in

chelates form, not ionic. Secondly, the role of other compound than chlorophyll in chloroplast that have antioxidant capacity, for example exogenous superoxide dismutation, especially Fe-SOD, Cu, Zn-SOD, Mn-SOD and ascorbic acid. Superoxide dismutase is an antioxidant that catalyzes superoxide anion superoxide (O[•]) and acts as the first defense against free radicals, whereas ascorbic acid breaks the peroxide radicals chain by giving hydrogen ions to free radicals and change into more stable molecules. And ascorbic acid helps superoxide dismutation role by catalyzing superoxide radicals reaction into H₂O₂ that more stable.²⁵ Thirdly, papaya leaves contain saponin, which also have antioxidant capacity. Saponin activates catalase and superoxide dismutation by giving hydrogen ions.²⁶

Combination of some antioxidant components in papaya leaves extract, which are chlorophyll, magnesium, superoxide dismutation exogenous, vitamin C, and saponins, have no possible prooxidant effects in experimental animals. Hematic components are prooxidant, while the structure of porphyrins in chlorophyll are antioxidants.²⁷ Giving oral treatment is considered to be safer than any other way. Saponins have toxic effects when given intravenously, but toxic effects are reduced when administered orally.²⁸

The effect of papaya leaf chlorophyll extract on blood glucose level. Both treatment were able to lower blood glucose levels close to normal (50-125 mg/dL) but the best effect occurred in treatment B, although it was not as good as normal rats without induction (technique control). The role of hypoglycemic agent from the extract is thought to be caused by 2 factors. Firstly, the role of some antioxidant components which are chlorophyll, magnesium, exogenous superoxide dismutation, and vitamin C in counteracting free radicals. Secondly, saponin content in papaya leaves also has a role as a hypoglycemic agent. Even saponin content in 22 Chinese herbs has antioxidant activity and high antiglycation, thus it plays a role in the treatment of diabetes mellitus.²⁹

Di Naso³⁰ also found that exogenous administration of superoxide dismutation had no effect on blood glucose levels in diabetic rats group. Exogenous superoxide dismutase in this study was orgotein (generic name Cu-Zn SOD), a pure substance derived from cow liver after going through the process of heating, enzymatic digestion with other proteins, as well as the purification of the homogenate by the method of ionic exchange chromatography. The absence of hypoglycemic effects was thought to be caused by the time and duration of exogenous superoxide dismutase administration. Although there were no effect on blood glucose levels, oxidative stress decreased, and there was increased antioxidant enzymatic activity in the liver of diabetic rats.

Conclusion

Extract of chlorophyll in papaya leaves for 40 days increased the antioxidative status of the body, indicated with low levels of superoxide dismutase and decreased blood glucose level. The higher dose of extract, the lower level of superoxide dismutase and the higher change of blood glucose level. The extract with a dose of 200.6 mg/200 g body weight can lower blood glucose level to normal after 40 days.

Acknowledgments

We thank the technician at PAU Center for the Study of Food and Nutrition, Gadjah Mada University. We also thank the National Education Ministry for their financial support through Beasiswa Unggulan Kementerian Pendidikan Nasional year 2010.

References

- Setiawan B, Suhartono E. Stres oksidatif dan peran antioksidan pada diabetes mellitus. *Majalah Kedokteran Indonesia*. 2005;55:86-91. [In Indonesia]
- Prangdimurti E, Muchtadi D, Astawan M, Zakaria FR. Aktivitas antioksidan ekstrak daun suji (*Pleomele angustifolia* NE Brown). *Jurnal Teknologi dan Industri Pangan* 2006;17:79-86.
- Endo Y, Usuki R, Kaneda T. Antioxidant effect of chlorophyll and pheophytin on the autooxidation of oils in the dark II. The mechanism of antioxidative action of chlorophyll. *JAOS*. 1985;62:1387-1388.
- Winarsi H. *Antioksidan alami dan radikal bebas: Potensi dan aplikasinya dalam kesehatan*. Yogyakarta: Kanisius; 2007. p.278.
- Shao HB, Chu LY, Lu ZH, Kang CM. Primary antioxidant free radical scavenging and redox signaling pathways in higher plant cells. *Int J Biol Sci*. 2008;4:8-14.
- Setiari N, Nurchayati Y. Eksplorasi kandungan klorofil pada beberapa sayuran hijau sebagai alternative bahan dasar food supplement. *BIOMA*. 2009;11:6-10.
- Ayoola PB, Adeyeye A. Phytochemical and nutrient evaluation of *Carica papaya* (papaw) leaves. *IJRRS*. 2010;5(3):325-328.
- Fakeye TO, Oladipupo T, Showande O, Ogunremi Y. Effect of coadministration of extract of *Carica papaya* Linn (family Caricaceae) on activity of two oral hypoglycemic agents. *Trop J Pharm Res*. 2007;6:671-678.
- Gosh MN. *Fundamentals of experimental pharmacology*. Calcutta: Scientific Book Agency; 1971. p.287.
- Braunwald E, Hauser S, Jameson J, India E. 17th ed. *Harrison's principles of internal medicine*. US: McGraw-Hill Companies, Inc; 2008. p.2958
- Youth RA, Simmerman SJ, Newell R, King RA. Ketamine anesthesia for rats. *Physiol Behav*. 2003; 10:633-636.
- Marklund S, Marklund G. Involvement of the superoxide anion radical in the autooxidation of pyrogallol and a convenient assay for superoxide dismutase. *Eur J Biochem*. 1974;47:469-474.
- Johnson-Delaney C. Exotic animal companion medicine handbook for veterinarians. 1996. Zoological educational network. In: Braslasu ED, Bradatan C, Cornila M, Savulescu I, Cojmaleata R, Braslasu MC. Normal blood glucose in white wistar rat and its changes following anesthesia. *Lucari Stiintifice Medicina Veterinara* 2007;40:120-123.
- Bond O. *The recommended daily dosages of chlorophyll*. (online) [cited 2011]. Available from: <http://www.livestrong.com/article/454170-the-recommended-daily-dosages-of-chlorophyll/>.
- Zafar M, Naeem-Ul-Hassan Naqvi S. Effect of stz-induced diabetes on the relative weights of kidney, liver and pancreas in albino rats: a comparative study. *Int J Morphol*. 2010;28:135-142.
- Takada J, Machado MA, Peres SB, Brito LC, Borges-Silva CN, Costa CEM, Fonseca-Alaniz MH, Andreotti S, Lima FB. Neonatal streptozotocin-induced diabetes mellitus: a model of insulin resistance associated with loss of adipose mass. *Metabolism*. 2007;56:977-984.
- Pepato MT, Migliorini RH, Goldberg AL, Kettelhud IC. Role of different proteolytic pathways in degradation of muscle protein from streptozotocin-induced rats. *Am J Physiol*. 1996;271:E340-E347.
- Zulfikar. *Peptida sebagai rantai protein*. (online) [cited 2010]. Available from: <http://www.chem-is-try.org>.
- Ventura-Sobrevilla J, Boone-Villa VD, Agullar CN, Roman-Ramos R, Vega-Avilla E, Campos-Sepulveda E, Alarcon-Agullar F. Effect of varying dose and administration of streptozotocin on blood sugar in male CD1 mice. *Proc West Pharmacol Soc*. 2011;54:5-9.
- Kakkar R, Mantha SV, Radhi J, Prasad K, Kalra J. Increased oxidative stress in rat liver and pancreas during progression of streptozotocin-induced diabetes. *Clin Sci*. 1998;94:623-632.
- Ellah MRA, Okada K, Goryo M, Oishi A, Yasuda J. Superoxide dismutase activity as a measure of hepatic oxidative stress in cattle following ethionine administration. *Vet. J*. 2009;182:336-341.
- Siburian M. *Aktivitas antioksidan superoksida dismutase pada hati tikus hiperkolesterolemia yang diberi ekstrak kulit mahoni (Swietenia macrophylla)* [Skripsi]. Bogor: Institut Pertanian Bogor; 2011. [In Indonesia]
- Choi J, Roche E, Caquet T. Characterization of superoxide dismutase activity in *Chironomus riparius* Mg. (Diptera, Chironomidae) larvae—a potential biomarker. *Comp Biochem Physiol Part C*. 1999;124:73-81.
- Moussa SA. Oxidative stress in diabetes mellitus. *Romanian J Biophys*. 2008;18:225-236.
- Wresdiyati T, Astawan M, Fithriani D, Adnyane IKM, Novelina S, Aryani S. Pengaruh α -tokoferol terhadap profil superoksida dismutase dan malondialdehid pada jaringan hati tikus di bawah kondisi stress. *Jurnal Veteriner*. 2002;13:202-209.
- Elekofehinti OO, Adanlawo IG, Fakoya A, Saliu JA, Sodehinde SA. Effects of saponin from *Solanum anguivi* Lam fruit on heart and kidney superoxide dismutase, catalase, and malondialdehyde in rat. *J Biol Sci*. 2012;4:530-533.
- Masushita S, Iwami N. Antioxidative abilities of some porphyrins on the oxidation of sodium linoleate. *Arch Biochem Biophys*. 1965;112:476-477.

28. Well B. Free radical and metal ions in health and disease. *Proc Nutr Soc.* 1987;46:13-26.
29. Chen HY, Yen GC. Antioxidant activity and free radical-scavenging capacity of extract from guava (*Psidium guajava* L) leaves. *J Food Chem.* 2006. In: Wang TS, editor. Relation between antioxidant and antiglycation ability of saponin, polyphenols, and polysaccharides in Chinese herbal medicines used to treat diabetes. *J Medicinal Plant Res.* 2011;5:2322-2331.
30. Di Naso FC, Simoes Dias A, Porawski M, Marroni NA. Exogenous superoxide dismutase: action on liver oxidative stress in animals with streptozotocin-induced diabetes. *Exp Diabetes Res.* 2011;2011:754132.