



Coronary Disease Risk Prediction Score Sheet for Men Based on LDL Cholesterol Level

Step 1

| Age | |
|-------|--------|
| Years | Points |
| 30-34 | -1 |
| 35-39 | 0 |
| 40-44 | 1 |
| 45-49 | 2 |
| 50-54 | 3 |
| 55-59 | 4 |
| 60-64 | 5 |
| 65-69 | 6 |
| 70-74 | 7 |

Step 2

| LDL - Cholesterol | | |
|-------------------|-----------|--------|
| (mg/dl) | (mmol/L) | Points |
| <100 | ≤2.59 | -3 |
| 100-129 | 2.60-3.36 | 0 |
| 130-159 | 3.37-4.14 | 0 |
| 160-189 | 4.15-4.91 | 1 |
| ≥190 | ≥4.92 | 2 |

| Key | |
|--------|-----------|
| Color | Risk |
| green | Very low |
| white | Low |
| yellow | Moderate |
| rose | High |
| red | Very high |

Step 3

| HDL - Cholesterol | | |
|-------------------|-----------|--------|
| (mg/dl) | (mmol/L) | Points |
| <35 | ≤0.90 | 2 |
| 35-44 | 0.91-1.16 | 1 |
| 45-49 | 1.17-1.29 | 0 |
| 50-59 | 1.30-1.55 | 0 |
| ≥60 | ≥1.56 | -1 |

Step 4

| Blood Pressure | | | | | |
|-----------------|------------------|-------|-------|-------|-------|
| Systolic (mmHg) | Diastolic (mmHg) | | | | |
| | <80 | 80-84 | 85-89 | 90-99 | ≥100 |
| <120 | 0 | | | | |
| 120-129 | | 0 pts | | | |
| 130-139 | | | 1 | | |
| 140-159 | | | | 2 | |
| ≥160 | | | | | 3 pts |

Note: When systolic and diastolic pressures provide different estimates for point scores, use the higher number

Step 5

| Diabetes | |
|----------|--------|
| | Points |
| No | 0 |
| Yes | 2 |

Step 6

| Smoker | |
|--------|--------|
| | Points |
| No | 0 |
| Yes | 2 |

Risk estimates were derived from the experience of the NHLBI's Framingham Heart Study, a predominantly Caucasian population in Massachusetts, USA

Step 7 (sum from steps 1-6)

| Adding up the points | |
|----------------------|-------|
| Age | _____ |
| LDL Cholesterol | _____ |
| HDL Cholesterol | _____ |
| Blood Pressure | _____ |
| Diabetes | _____ |
| Smoker | _____ |
| Point Total | _____ |

Step 8 (determine CHD risk from point total)

| CHD Risk | |
|-------------|----------------|
| Point Total | 10 Yr CHD Risk |
| ≤-3 | 1% |
| -2 | 2% |
| -1 | 2% |
| 0 | 3% |
| 1 | 4% |
| 2 | 4% |
| 3 | 6% |
| 4 | 7% |
| 5 | 9% |
| 6 | 11% |
| 7 | 14% |
| 8 | 18% |
| 9 | 22% |
| 10 | 27% |
| 11 | 33% |
| 12 | 40% |
| 13 | 47% |
| ≥14 | ≥56% |

Step 9 (compare to man of the same age)

| Age (years) | Comparative Risk | |
|-------------|------------------------|---------------------|
| | Average 10 Yr CHD Risk | Low* 10 Yr CHD Risk |
| 30-34 | 3% | 2% |
| 35-39 | 5% | 3% |
| 40-44 | 7% | 4% |
| 45-49 | 11% | 4% |
| 50-54 | 14% | 6% |
| 55-59 | 16% | 7% |
| 60-64 | 21% | 9% |
| 65-69 | 25% | 11% |
| 70-74 | 30% | 14% |

*Low risk was calculated for a man the same age, normal blood pressure, LDL cholesterol 100-129 mg/dL, HDL cholesterol 45 mg/dL, non-smoker, no diabetes

Analisis Univariat

Tingkat Risiko PJK

| | Frequency | Percent | Valid Percent | Cumulative Percent |
|--------------|-----------|---------|---------------|--------------------|
| Valid rendah | 79 | 34.1 | 34.1 | 34.1 |
| sedang | 134 | 57.8 | 57.8 | 91.8 |
| tinggi | 19 | 8.2 | 8.2 | 100.0 |
| Total | 232 | 100.0 | 100.0 | |

Kadar Kolesterol Total

| | Frequency | Percent | Valid Percent | Cumulative Percent |
|------------|-----------|---------|---------------|--------------------|
| Valid <200 | 99 | 42.7 | 42.7 | 42.7 |
| >= 200 | 133 | 57.3 | 57.3 | 100.0 |
| Total | 232 | 100.0 | 100.0 | |

Kadar LDL

| | Frequency | Percent | Valid Percent | Cumulative Percent |
|------------|-----------|---------|---------------|--------------------|
| Valid <130 | 112 | 48.3 | 48.3 | 48.3 |
| >= 130 | 120 | 51.7 | 51.7 | 100.0 |
| Total | 232 | 100.0 | 100.0 | |

Kadar HDL

| | Frequency | Percent | Valid Percent | Cumulative Percent |
|-----------|-----------|---------|---------------|--------------------|
| Valid <40 | 67 | 28.9 | 28.9 | 28.9 |
| >=40 | 165 | 71.1 | 71.1 | 100.0 |
| Total | 232 | 100.0 | 100.0 | |

Kadar gula darah puasa

| | Frequency | Percent | Valid Percent | Cumulative Percent |
|------------|-----------|---------|---------------|--------------------|
| Valid <110 | 189 | 81.5 | 81.5 | 81.5 |
| >=110 | 43 | 18.5 | 18.5 | 100.0 |
| Total | 232 | 100.0 | 100.0 | |

IMT

| | Frequency | Percent | Valid Percent | Cumulative Percent |
|-----------|-----------|---------|---------------|--------------------|
| Valid <25 | 109 | 47.0 | 47.0 | 47.0 |
| >=25 | 123 | 53.0 | 53.0 | 100.0 |
| Total | 232 | 100.0 | 100.0 | |

Usia Karyawan

| | Frequency | Percent | Valid Percent | Cumulative Percent |
|-----------------|-----------|---------|---------------|--------------------|
| Valid <50 tahun | 137 | 59.1 | 59.1 | 59.1 |
| >= 50 tahun | 95 | 40.9 | 40.9 | 100.0 |
| Total | 232 | 100.0 | 100.0 | |

Tekanan Darah Sistolik dan Diastolik

| | Frequency | Percent | Valid Percent | Cumulative Percent |
|---------------|-----------|---------|---------------|--------------------|
| Valid <120/80 | 150 | 64.7 | 64.7 | 64.7 |
| >=120/80 | 82 | 35.3 | 35.3 | 100.0 |
| Total | 232 | 100.0 | 100.0 | |

Status Merokok

| | Frequency | Percent | Valid Percent | Cumulative Percent |
|---------------------|-----------|---------|---------------|--------------------|
| Valid tidak merokok | 126 | 54.3 | 54.3 | 54.3 |
| merokok | 106 | 45.7 | 45.7 | 100.0 |
| Total | 232 | 100.0 | 100.0 | |

Jabatan Karyawan

| | Frequency | Percent | Valid Percent | Cumulative Percent |
|----------------------|-----------|---------|---------------|--------------------|
| Valid jabatan rendah | 207 | 89.2 | 89.2 | 89.2 |
| jabatan tinggi | 25 | 10.8 | 10.8 | 100.0 |
| Total | 232 | 100.0 | 100.0 | |

Kategori kolesterol Total * Kategori Risk PJK Crosstabulation

| | | | Kategori Risk PJK | | | Total |
|---------------------------|------------------------------------|------------------------------------|-------------------|--------|--------|--------|
| | | | rendah | sedang | tinggi | |
| Kategori kolesterol Total | <200 | Count | 43 | 52 | 4 | 99 |
| | | % within Kategori kolesterol Total | 43.4% | 52.5% | 4.0% | 100.0% |
| | ≥ 200 | Count | 36 | 82 | 15 | 133 |
| | | % within Kategori kolesterol Total | 27.1% | 61.7% | 11.3% | 100.0% |
| Total | Count | 79 | 134 | 19 | 232 | |
| | % within Kategori kolesterol Total | 34.1% | 57.8% | 8.2% | 100.0% | |

Kategori LDL * Kategori Risk PJK Crosstabulation

| | | | Kategori Risk PJK | | | Total |
|--------------|-----------------------|-----------------------|-------------------|--------|--------|--------|
| | | | rendah | sedang | tinggi | |
| Kategori LDL | <130 | Count | 50 | 56 | 6 | 112 |
| | | % within Kategori LDL | 44.6% | 50.0% | 5.4% | 100.0% |
| | ≥ 130 | Count | 29 | 78 | 13 | 120 |
| | | % within Kategori LDL | 24.2% | 65.0% | 10.8% | 100.0% |
| Total | Count | 79 | 134 | 19 | 232 | |
| | % within Kategori LDL | 34.1% | 57.8% | 8.2% | 100.0% | |

Kategori HDL * Kategori Risk PJK Crosstabulation

| | | | Kategori Risk PJK | | | Total |
|--------------|-----------------------|-----------------------|-------------------|--------|--------|--------|
| | | | rendah | sedang | tinggi | |
| Kategori HDL | <40 | Count | 18 | 38 | 11 | 67 |
| | | % within Kategori HDL | 26.9% | 56.7% | 16.4% | 100.0% |
| | ≥40 | Count | 61 | 96 | 8 | 165 |
| | | % within Kategori HDL | 37.0% | 58.2% | 4.8% | 100.0% |
| Total | Count | 79 | 134 | 19 | 232 | |
| | % within Kategori HDL | 34.1% | 57.8% | 8.2% | 100.0% | |

Kategori kadar gula * Kategori Risk PJK Crosstabulation

| | | | Kategori Risk PJK | | | Total |
|---------------------|-------|------------------------------|-------------------|--------|--------|--------|
| | | | rendah | sedang | tinggi | |
| Kategori kadar gula | <110 | Count | 75 | 106 | 8 | 189 |
| | | % within Kategori kadar gula | 39.7% | 56.1% | 4.2% | 100.0% |
| | >=110 | Count | 4 | 28 | 11 | 43 |
| | | % within Kategori kadar gula | 9.3% | 65.1% | 25.6% | 100.0% |
| Total | | Count | 79 | 134 | 19 | 232 |
| | | % within Kategori kadar gula | 34.1% | 57.8% | 8.2% | 100.0% |

Kategori IMT * Kategori Risk PJK Crosstabulation

| | | | Kategori Risk PJK | | | Total |
|--------------|------|-----------------------|-------------------|--------|--------|--------|
| | | | rendah | sedang | tinggi | |
| Kategori IMT | <25 | Count | 37 | 66 | 6 | 109 |
| | | % within Kategori IMT | 33.9% | 60.6% | 5.5% | 100.0% |
| | >=25 | Count | 42 | 68 | 13 | 123 |
| | | % within Kategori IMT | 34.1% | 55.3% | 10.6% | 100.0% |
| Total | | Count | 79 | 134 | 19 | 232 |
| | | % within Kategori IMT | 34.1% | 57.8% | 8.2% | 100.0% |

usia_Pkj * Kategori Risk PJK Crosstabulation

| | | | Kategori Risk PJK | | | Total |
|----------|-------------|-------------------|-------------------|--------|--------|--------|
| | | | rendah | sedang | tinggi | |
| usia_Pkj | <50 tahun | Count | 37 | 90 | 10 | 137 |
| | | % within usia_Pkj | 27.0% | 65.7% | 7.3% | 100.0% |
| | >= 50 tahun | Count | 42 | 44 | 9 | 95 |
| | | % within usia_Pkj | 44.2% | 46.3% | 9.5% | 100.0% |
| Total | | Count | 79 | 134 | 19 | 232 |
| | | % within usia_Pkj | 34.1% | 57.8% | 8.2% | 100.0% |

Tek_darah * Kategori Risk PJK Crosstabulation

| | | | Kategori Risk PJK | | | Total |
|-----------|----------|--------------------|-------------------|--------|--------|--------|
| | | | rendah | sedang | tinggi | |
| Tek_darah | <120/80 | Count | 60 | 83 | 7 | 150 |
| | | % within Tek_darah | 40.0% | 55.3% | 4.7% | 100.0% |
| | >=120/80 | Count | 19 | 51 | 12 | 82 |
| | | % within Tek_darah | 23.2% | 62.2% | 14.6% | 100.0% |
| Total | | Count | 79 | 134 | 19 | 232 |
| | | % within Tek_darah | 34.1% | 57.8% | 8.2% | 100.0% |

Merokok * Kategori Risk PJK Crosstabulation

| | | | Kategori Risk PJK | | | Total |
|---------|---------------|------------------|-------------------|--------|--------|--------|
| | | | rendah | sedang | tinggi | |
| Merokok | tidak merokok | Count | 65 | 59 | 2 | 126 |
| | | % within Merokok | 51.6% | 46.8% | 1.6% | 100.0% |
| | merokok | Count | 14 | 75 | 17 | 106 |
| | | % within Merokok | 13.2% | 70.8% | 16.0% | 100.0% |
| Total | | Count | 79 | 134 | 19 | 232 |
| | | % within Merokok | 34.1% | 57.8% | 8.2% | 100.0% |

Jabatan * Kategori Risk PJK Crosstabulation

| | | | Kategori Risk PJK | | | Total |
|---------|----------------|------------------|-------------------|--------|--------|--------|
| | | | rendah | sedang | tinggi | |
| Jabatan | jabatan rendah | Count | 69 | 119 | 19 | 207 |
| | | % within Jabatan | 33.3% | 57.5% | 9.2% | 100.0% |
| | jabatan tinggi | Count | 10 | 15 | 0 | 25 |
| | | % within Jabatan | 40.0% | 60.0% | .0% | 100.0% |
| Total | | Count | 79 | 134 | 19 | 232 |
| | | % within Jabatan | 34.1% | 57.8% | 8.2% | 100.0% |

Analisis Bivariat

Crosstab

| | | | Risk_PJK1 | | Total |
|---------------------------|-------|------------------------------------|---------------|---------------|--------|
| | | | risiko rendah | risiko tinggi | |
| Kategori kolesterol Total | <200 | Count | 43 | 56 | 99 |
| | | % within Kategori kolesterol Total | 43.4% | 56.6% | 100.0% |
| | >=200 | Count | 36 | 97 | 133 |
| | | % within Kategori kolesterol Total | 27.1% | 72.9% | 100.0% |
| Total | | Count | 79 | 153 | 232 |
| | | % within Kategori kolesterol Total | 34.1% | 65.9% | 100.0% |

Chi-Square Tests

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|------------------------------------|--------------------|----|-----------------------|----------------------|----------------------|
| Pearson Chi-Square | 6.770 ^b | 1 | .009 | | |
| Continuity Correction ^a | 6.061 | 1 | .014 | | |
| Likelihood Ratio | 6.743 | 1 | .009 | | |
| Fisher's Exact Test | | | | .012 | .007 |
| Linear-by-Linear Association | 6.741 | 1 | .009 | | |
| N of Valid Cases | 232 | | | | |

a. Computed only for a 2x2 table

b. 0 cells (.0%) have expected count less than 5. The minimum expected count is 33.71.

Risk Estimate

| | Value | 95% Confidence Interval | |
|---------------------------------------------------------|-------|-------------------------|-------|
| | | Lower | Upper |
| Odds Ratio for Kategori kolesterol Total (<200 / >=200) | 2.069 | 1.192 | 3.592 |
| For cohort Risk_PJK1 = risiko rendah | 1.605 | 1.121 | 2.296 |
| For cohort Risk_PJK1 = risiko tinggi | .776 | .634 | .949 |
| N of Valid Cases | 232 | | |

Crosstab

| | | | Risk_PJK1 | | Total |
|--------------|-------|-----------------------|---------------|---------------|--------|
| | | | risiko rendah | risiko tinggi | |
| Kategori LDL | <130 | Count | 50 | 62 | 112 |
| | | % within Kategori LDL | 44.6% | 55.4% | 100.0% |
| | >=130 | Count | 29 | 91 | 120 |
| | | % within Kategori LDL | 24.2% | 75.8% | 100.0% |
| Total | | Count | 79 | 153 | 232 |
| | | % within Kategori LDL | 34.1% | 65.9% | 100.0% |

Chi-Square Tests

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|------------------------------------|---------------------|----|-----------------------|----------------------|----------------------|
| Pearson Chi-Square | 10.816 ^b | 1 | .001 | | |
| Continuity Correction ^a | 9.923 | 1 | .002 | | |
| Likelihood Ratio | 10.904 | 1 | .001 | | |
| Fisher's Exact Test | | | | .001 | .001 |
| Linear-by-Linear Association | 10.769 | 1 | .001 | | |
| N of Valid Cases | 232 | | | | |

a. Computed only for a 2x2 table

b. 0 cells (.0%) have expected count less than 5. The minimum expected count is 38.14.

Risk Estimate

| | Value | 95% Confidence Interval | |
|--------------------------------------------|-------|-------------------------|-------|
| | | Lower | Upper |
| Odds Ratio for Kategori LDL (<130 / >=130) | 2.531 | 1.446 | 4.430 |
| For cohort Risk_PJK1 = risiko rendah | 1.847 | 1.266 | 2.696 |
| For cohort Risk_PJK1 = risiko tinggi | .730 | .601 | .887 |
| N of Valid Cases | 232 | | |

Crosstab

| | | Risk_PJK1 | | Total | |
|--------------|-------|-----------------------|---------------|-------|--------|
| | | risiko rendah | risiko tinggi | | |
| Kategori HDL | <40 | Count | 18 | 49 | 67 |
| | | % within Kategori HDL | 26.9% | 73.1% | 100.0% |
| >=40 | Count | 61 | 104 | 165 | |
| | | % within Kategori HDL | 37.0% | 63.0% | 100.0% |
| Total | Count | 79 | 153 | 232 | |
| | | % within Kategori HDL | 34.1% | 65.9% | 100.0% |

Chi-Square Tests

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|------------------------------------|--------------------|----|-----------------------|----------------------|----------------------|
| Pearson Chi-Square | 2.166 ^b | 1 | .141 | | |
| Continuity Correction ^a | 1.740 | 1 | .187 | | |
| Likelihood Ratio | 2.220 | 1 | .136 | | |
| Fisher's Exact Test | | | | .169 | .093 |
| Linear-by-Linear Association | 2.157 | 1 | .142 | | |
| N of Valid Cases | 232 | | | | |

a. Computed only for a 2x2 table

b. 0 cells (.0%) have expected count less than 5. The minimum expected count is 22.81.

Risk Estimate

| | Value | 95% Confidence Interval | |
|------------------------------------------|-------|-------------------------|-------|
| | | Lower | Upper |
| Odds Ratio for Kategori HDL (<40 / >=40) | .626 | .335 | 1.171 |
| For cohort Risk_PJK1 = risiko rendah | .727 | .467 | 1.131 |
| For cohort Risk_PJK1 = risiko tinggi | 1.160 | .963 | 1.398 |
| N of Valid Cases | 232 | | |

Crosstab

| | | Risk_PJK1 | | Total |
|---------------------|-------|---------------------------------------|---------------------------------------|----------------------------------------|
| | | risiko rendah | risiko tinggi | |
| Kategori kadar gula | <110 | Count 75 | Count 114 | Count 189 |
| | | % within Kategori kadar gula 39.7% | % within Kategori kadar gula 60.3% | % within Kategori kadar gula 100.0% |
| | >=110 | Count 4 | Count 39 | Count 43 |
| | | % within Kategori kadar gula 9.3% | % within Kategori kadar gula 90.7% | % within Kategori kadar gula 100.0% |
| Total | | Count 79 | Count 153 | Count 232 |
| | | % within Kategori kadar gula 34.1% | % within Kategori kadar gula 65.9% | % within Kategori kadar gula 100.0% |

Chi-Square Tests

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|------------------------------------|---------------------|----|-----------------------|----------------------|----------------------|
| Pearson Chi-Square | 14.397 ^b | 1 | .000 | | |
| Continuity Correction ^a | 13.076 | 1 | .000 | | |
| Likelihood Ratio | 17.080 | 1 | .000 | | |
| Fisher's Exact Test | | | | .000 | .000 |
| Linear-by-Linear Association | 14.335 | 1 | .000 | | |
| N of Valid Cases | 232 | | | | |

a. Computed only for a 2x2 table

b. 0 cells (.0%) have expected count less than 5. The minimum expected count is 14.64.

Risk Estimate

| | Value | 95% Confidence Interval | |
|---------------------------------------------------|-------|-------------------------|--------|
| | | Lower | Upper |
| Odds Ratio for Kategori kadar gula (<110 / >=110) | 6.414 | 2.201 | 18.691 |
| For cohort Risk_PJK1 = risiko rendah | 4.266 | 1.650 | 11.027 |
| For cohort Risk_PJK1 = risiko tinggi | .665 | .572 | .773 |
| N of Valid Cases | 232 | | |

Crosstab

| | | | Risk_PJK1 | | Total |
|-----------------|------|-----------------------|---------------|---------------|--------|
| | | | risiko rendah | risiko tinggi | |
| Kategori IMT | <25 | Count | 37 | 72 | 109 |
| | | % within Kategori IMT | 33.9% | 66.1% | 100.0% |
| | >=25 | Count | 42 | 81 | 123 |
| | | % within Kategori IMT | 34.1% | 65.9% | 100.0% |
| Total | | Count | 79 | 153 | 232 |
| | | % within Kategori IMT | 34.1% | 65.9% | 100.0% |

Chi-Square Tests

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|------------------------------------|-------------------|----|--------------------------|-------------------------|-------------------------|
| Pearson Chi-Square | .001 ^b | 1 | .974 | | |
| Continuity Correction ^a | .000 | 1 | 1.000 | | |
| Likelihood Ratio | .001 | 1 | .974 | | |
| Fisher's Exact Test | | | | 1.000 | .543 |
| Linear-by-Linear Association | .001 | 1 | .974 | | |
| N of Valid Cases | 232 | | | | |

a. Computed only for a 2x2 table

b. 0 cells (.0%) have expected count less than 5. The minimum expected count is 37.12.

Risk Estimate

| | Value | 95% Confidence Interval | |
|---------------------------------------------|-------|-------------------------|-------|
| | | Lower | Upper |
| Odds Ratio for Kategori IMT (<25 / >=25) | .991 | .575 | 1.708 |
| For cohort Risk_PJK1 = risiko rendah | .994 | .694 | 1.423 |
| For cohort Risk_PJK1 = risiko tinggi | 1.003 | .833 | 1.207 |
| N of Valid Cases | 232 | | |

Crosstab

| | | | Risk_PJK1 | | Total |
|----------|------|-------------------|---------------|---------------|--------|
| | | | risiko rendah | risiko tinggi | |
| usia_Pkj | <50 | Count | 37 | 100 | 137 |
| | | % within usia_Pkj | 27.0% | 73.0% | 100.0% |
| | >=50 | Count | 42 | 53 | 95 |
| | | % within usia_Pkj | 44.2% | 55.8% | 100.0% |
| Total | | Count | 79 | 153 | 232 |
| | | % within usia_Pkj | 34.1% | 65.9% | 100.0% |

Chi-Square Tests

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|------------------------------------|--------------------|----|-----------------------|----------------------|----------------------|
| Pearson Chi-Square | 7.393 ^b | 1 | .007 | | |
| Continuity Correction ^a | 6.647 | 1 | .010 | | |
| Likelihood Ratio | 7.345 | 1 | .007 | | |
| Fisher's Exact Test | | | | .008 | .005 |
| Linear-by-Linear Association | 7.361 | 1 | .007 | | |
| N of Valid Cases | 232 | | | | |

a. Computed only for a 2x2 table

b. 0 cells (.0%) have expected count less than 5. The minimum expected count is 32.35.

Risk Estimate

| | Value | 95% Confidence Interval | |
|--------------------------------------|-------|-------------------------|-------|
| | | Lower | Upper |
| Odds Ratio for usia_Pkj (<50 / >=50) | .467 | .268 | .812 |
| For cohort Risk_PJK1 = risiko rendah | .611 | .428 | .872 |
| For cohort Risk_PJK1 = risiko tinggi | 1.308 | 1.065 | 1.608 |
| N of Valid Cases | 232 | | |

Crosstab

| | | Risk_PJK1 | | Total | |
|-----------|--------|--------------------|---------------|-------|--------|
| | | risiko rendah | risiko tinggi | | |
| Tek_darah | normal | Count | 60 | 90 | 150 |
| | | % within Tek_darah | 40.0% | 60.0% | 100.0% |
| tinggi | Count | 19 | 63 | 82 | |
| | | % within Tek_darah | 23.2% | 76.8% | 100.0% |
| Total | Count | 79 | 153 | 232 | |
| | | % within Tek_darah | 34.1% | 65.9% | 100.0% |

Chi-Square Tests

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|------------------------------------|--------------------|----|-----------------------|----------------------|----------------------|
| Pearson Chi-Square | 6.687 ^b | 1 | .010 | | |
| Continuity Correction ^a | 5.958 | 1 | .015 | | |
| Likelihood Ratio | 6.918 | 1 | .009 | | |
| Fisher's Exact Test | | | | .013 | .007 |
| Linear-by-Linear Association | 6.658 | 1 | .010 | | |
| N of Valid Cases | 232 | | | | |

a. Computed only for a 2x2 table

b. 0 cells (.0%) have expected count less than 5. The minimum expected count is 27.92.

Risk Estimate

| | Value | 95% Confidence Interval | |
|--------------------------------------------|-------|-------------------------|-------|
| | | Lower | Upper |
| Odds Ratio for Tek_darah (normal / tinggi) | 2.211 | 1.203 | 4.061 |
| For cohort Risk_PJK1 = risiko rendah | 1.726 | 1.112 | 2.681 |
| For cohort Risk_PJK1 = risiko tinggi | .781 | .655 | .932 |
| N of Valid Cases | 232 | | |

Crosstab

| | | Risk_PJK1 | | Total | |
|---------|-------|------------------|---------------|-------|--------|
| | | risiko rendah | risiko tinggi | | |
| Merokok | tidak | Count | 65 | 61 | 126 |
| | | % within Merokok | 51.6% | 48.4% | 100.0% |
| | ya | Count | 14 | 92 | 106 |
| | | % within Merokok | 13.2% | 86.8% | 100.0% |
| Total | | Count | 79 | 153 | 232 |
| | | % within Merokok | 34.1% | 65.9% | 100.0% |

Chi-Square Tests

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|------------------------------------|---------------------|----|-----------------------|----------------------|----------------------|
| Pearson Chi-Square | 37.762 ^b | 1 | .000 | | |
| Continuity Correction ^a | 36.072 | 1 | .000 | | |
| Likelihood Ratio | 40.307 | 1 | .000 | | |
| Fisher's Exact Test | | | | .000 | .000 |
| Linear-by-Linear Association | 37.599 | 1 | .000 | | |
| N of Valid Cases | 232 | | | | |

a. Computed only for a 2x2 table

b. 0 cells (.0%) have expected count less than 5. The minimum expected count is 36.09.

Risk Estimate

| | Value | 95% Confidence Interval | |
|--------------------------------------|-------|-------------------------|--------|
| | | Lower | Upper |
| Odds Ratio for Merokok (tidak / ya) | 7.002 | 3.612 | 13.575 |
| For cohort Risk_PJK1 = risiko rendah | 3.906 | 2.330 | 6.547 |
| For cohort Risk_PJK1 = risiko tinggi | .558 | .459 | .678 |
| N of Valid Cases | 232 | | |

Crosstab

| | | | Risk_PJK1 | | Total |
|---------|--------|------------------|---------------|---------------|--------|
| | | | risiko rendah | risiko tinggi | |
| Jabatan | rendah | Count | 69 | 138 | 207 |
| | | % within Jabatan | 33.3% | 66.7% | 100.0% |
| | tinggi | Count | 10 | 15 | 25 |
| | | % within Jabatan | 40.0% | 60.0% | 100.0% |
| Total | | Count | 79 | 153 | 232 |
| | | % within Jabatan | 34.1% | 65.9% | 100.0% |

Chi-Square Tests

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|------------------------------------|-------------------|----|-----------------------|----------------------|----------------------|
| Pearson Chi-Square | .441 ^b | 1 | .506 | .510 | .324 |
| Continuity Correction ^a | .195 | 1 | .659 | | |
| Likelihood Ratio | .432 | 1 | .511 | | |
| Fisher's Exact Test | | | | | |
| Linear-by-Linear Association | .440 | 1 | .507 | | |
| N of Valid Cases | 232 | | | | |

a. Computed only for a 2x2 table

b. 0 cells (.0%) have expected count less than 5. The minimum expected count is 8.51.

Risk Estimate

| | Value | 95% Confidence Interval | |
|------------------------------------------|-------|-------------------------|-------|
| | | Lower | Upper |
| Odds Ratio for Jabatan (rendah / tinggi) | .750 | .320 | 1.756 |
| For cohort Risk_PJK1 = risiko rendah | .833 | .497 | 1.398 |
| For cohort Risk_PJK1 = risiko tinggi | 1.111 | .795 | 1.552 |
| N of Valid Cases | 232 | | |