

Statistics

Usia

N	Valid	30
	Missing	0

Usia

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid <= 60 Tahun	6	20.0	20.0	20.0
> 60 Tahun	24	80.0	80.0	100.0
Total	30	100.0	100.0	

Statistics

Jenis_Kelamin

N	Valid	30
	Missing	0

Jenis_Kelamin

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Laki-laki	7	23.3	23.3	23.3
Perempuan	23	76.7	76.7	100.0
Total	30	100.0	100.0	

Statistics

Pendidikan

N	Valid	30
	Missing	0

Pendidikan

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid S1	25	83.3	83.3	83.3
S2	5	16.7	16.7	100.0
Total	30	100.0	100.0	

Statistics

Lama_Kerjasam

N	Valid	30
	Missing	0

Lama_Kerjasam

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	<=5 tahun	8	26.7	26.7	26.7
	> 5 tahun	22	73.3	73.3	100.0
	Total	30	100.0	100.0	

Statistics

Pelatihan

N	Valid	30
	Missing	0

Pelatihan

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	pernah ikut	7	23.3	23.3	23.3
	tidak pernah ikut	23	76.7	76.7	100.0
	Total	30	100.0	100.0	

Statistics

pengetahuandokter

N	Valid	30
	Missing	0

pengetahuandokter

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	tidak tahu	8	26.7	26.7	26.7
	tahu	22	73.3	73.3	100.0
	Total	30	100.0	100.0	

Statistics

tahu1

N	Valid	30
	Missing	0

tahu1

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid tahu	30	100.0	100.0	100.0

Statistics

tahu2

N	Valid	30
	Missing	0

tahu2

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid tidak tahu	1	3.3	3.3	3.3
tahu	29	96.7	96.7	100.0
Total	30	100.0	100.0	

Statistics

tahu3

N	Valid	30
	Missing	0

tahu3

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid tidak tahu	8	26.7	26.7	26.7
tahu	22	73.3	73.3	100.0
Total	30	100.0	100.0	

Statistics

tahu4

N	Valid	30
	Missing	0

tahu4

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	tidak tahu	9	30.0	30.0	30.0
	tahu	21	70.0	70.0	100.0
	Total	30	100.0	100.0	

Statistics

tahu5

N	Valid	30
	Missing	0

tahu5

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	tidak tahu	5	16.7	16.7	16.7
	tahu	25	83.3	83.3	100.0
	Total	30	100.0	100.0	

Statistics

tahu6

N	Valid	30
	Missing	0

tahu6

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	tidak tahu	4	13.3	13.3	13.3
	tahu	26	86.7	86.7	100.0
	Total	30	100.0	100.0	

Statistics

tahu7

N	Valid	30
	Missing	0

tahu7

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	tidak tahu	7	23.3	23.3	23.3
	tahu	23	76.7	76.7	100.0
	Total	30	100.0	100.0	

Statistics

puas1

N	Valid	30
	Missing	0

puas1

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	tidak puas	21	70.0	70.0	70.0
	puas	9	30.0	30.0	100.0
	Total	30	100.0	100.0	

Statistics

puas2

N	Valid	30
	Missing	0

puas2

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	tidak puas	20	66.7	66.7	66.7
	puas	10	33.3	33.3	100.0
	Total	30	100.0	100.0	

kepuasandokter

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	tidak puas	20	66.7	66.7	66.7
	puas	10	33.3	33.3	100.0
	Total	30	100.0	100.0	

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	tidak tahu	8	26.7	26.7	26.7
	tahu	22	73.3	73.3	100.0
	Total	30	100.0	100.0	

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Usia * kepuasandokter	30	100.0%	0	.0%	30	100.0%

Usia * kepuasandokter Crosstabulation

		kepuasandokter		Total	
		tidak puas	puas		
Usia	<= 60 Tahun	Count	5	1	6
		% within Usia	83.3%	16.7%	100.0%
> 60 Tahun	Count	15	9	24	
		% within Usia	62.5%	37.5%	100.0%
Total	Count	20	10	30	
		% within Usia	66.7%	33.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.938(b)	1	.333		
Continuity Correction(a)	.234	1	.628		
Likelihood Ratio	1.029	1	.310		
Fisher's Exact Test				.633	.326
Linear-by-Linear Association	.906	1	.341		
N of Valid Cases	30				

a Computed only for a 2x2 table

b 2 cells (50.0%) have expected count less than 5. The minimum expected count is 2.00.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Usia (<= 60 Tahun / > 60 Tahun)	3.000	.301	29.940
For cohort kepuasandokter = tidak puas	1.333	.831	2.141
For cohort kepuasandokter = puas	.444	.069	2.861
N of Valid Cases	30		

Crosstabs

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Jenis_Kelamin * kepuasandokter	30	100.0%	0	.0%	30	100.0%

			kepuasandokter		Total
			tidak puas	puas	
Jenis_Kelamin	Laki-laki	Count	2	5	7
		% within Jenis_Kelamin	28.6%	71.4%	100.0%
	Perempuan	Count	18	5	23
		% within Jenis_Kelamin	78.3%	21.7%	100.0%
Total		Count	20	10	30
		% within Jenis_Kelamin	66.7%	33.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	5.963(b)	1	.015		
Continuity Correction(a)	3.936	1	.047		
Likelihood Ratio	5.730	1	.017		
Fisher's Exact Test				.026	.026
Linear-by-Linear Association	5.764	1	.016		
N of Valid Cases	30				

a Computed only for a 2x2 table

b 2 cells (50.0%) have expected count less than 5. The minimum expected count is 2.33.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Pendidikan * kepuasandokter	30	100.0%	0	.0%	30	100.0%

Pendidikan * kepuasandokter Crosstabulation

			kepuasandokter		Total
			tidak puas	puas	
Pendidikan	S1	Count	15	10	25
		% within Pendidikan	60.0%	40.0%	100.0%
	S2	Count	5	0	5
		% within Pendidikan	100.0%	.0%	100.0%
Total		Count	20	10	30
		% within Pendidikan	66.7%	33.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	3.000(b)	1	.083		
Continuity Correction(a)	1.470	1	.225		
Likelihood Ratio	4.540	1	.033		
Fisher's Exact Test				.140	.109
Linear-by-Linear Association	2.900	1	.089		
N of Valid Cases	30				

a Computed only for a 2x2 table

b 2 cells (50.0%) have expected count less than 5. The minimum expected count is 1.67.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
For cohort kepuasandokter = tidak puas	.600	.436	.826
N of Valid Cases	30		

Lama kerjasama*kepuasan dokter

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Lama_Kerjasam * kepuasandokter	30	100.0%	0	.0%	30	100.0%

			kepuasandokter		Total
			tidak puas	puas	
Lama_Kerjasam	<=5 tahun	Count	5	3	8
		% within Lama_Kerjasam	62.5%	37.5%	100.0%
	> 5 tahun	Count	15	7	22
		% within Lama_Kerjasam	68.2%	31.8%	100.0%
Total		Count	20	10	30
		% within Lama_Kerjasam	66.7%	33.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.085(b)	1	.770		
Continuity Correction(a)	.000	1	1.000		
Likelihood Ratio	.084	1	.772		
Fisher's Exact Test				1.000	.548
Linear-by-Linear Association	.082	1	.774		
N of Valid Cases	30				

a Computed only for a 2x2 table

b 1 cells (25.0%) have expected count less than 5. The minimum expected count is 2.67.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Lama_Kerjasam (<=5 tahun / > 5 tahun)	.778	.144	4.212
For cohort kepuasandokter = tidak puas	.917	.499	1.684
For cohort kepuasandokter = puas	1.179	.399	3.483
N of Valid Cases	30		

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Pelatihan * kepuasandokter	30	100.0%	0	.0%	30	100.0%

Pelatihan * kepuasandokter Crosstabulation

			kepuasandokter		Total
			tidak puas	puas	
Pelatihan	pernah ikut	Count	2	5	7
		% within Pelatihan	28.6%	71.4%	100.0%
	tidak pernah ikut	Count	18	5	23
		% within Pelatihan	78.3%	21.7%	100.0%
Total		Count	20	10	30
		% within Pelatihan	66.7%	33.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	5.963(b)	1	.015		
Continuity Correction(a)	3.936	1	.047		
Likelihood Ratio	5.730	1	.017		
Fisher's Exact Test				.026	.026
Linear-by-Linear Association	5.764	1	.016		
N of Valid Cases	30				

a Computed only for a 2x2 table

b 2 cells (50.0%) have expected count less than 5. The minimum expected count is 2.33.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Pelatihan (pernah ikut / tidak pernah ikut)	.111	.016	.755
For cohort kepuasandokter = tidak puas	.365	.111	1.201
For cohort kepuasandokter = puas	3.286	1.328	8.130
N of Valid Cases	30		

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
pengetahuandokter * kepuasandokter	30	100.0%	0	.0%	30	100.0%

pengetahuandokter * kepuasandokter Crosstabulation

			kepuasandokter		Total
			tidak puas	puas	
pengetahuandokter	tidak tahu	Count	8	0	8
		% within pengetahuandokter	100.0%	.0%	100.0%
	tahu	Count	12	10	22
		% within pengetahuandokter	54.5%	45.5%	100.0%
Total		Count	20	10	30
		% within pengetahuandokter	66.7%	33.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	5.455(b)	1	.020		
Continuity Correction(a)	3.601	1	.058		
Likelihood Ratio	7.874	1	.005		
Fisher's Exact Test				.029	.022
Linear-by-Linear Association	5.273	1	.022		
N of Valid Cases	30				

a Computed only for a 2x2 table

b 1 cells (25.0%) have expected count less than 5. The minimum expected count is 2.67.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
For cohort kepuasandokter = tidak puas	1.833	1.252	2.685
N of Valid Cases	30		

Warnings

No measures of association are computed for the crosstabulation of tahuB * kepuasandokter. At least one variable in each 2-way table upon which measures of association are computed is a constant.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
tahuB * kepuasandokter	30	100,0%	0	,0%	30	100,0%

tahuB * kepuasandokter Crosstabulation

			kepuasandokter		Total
			tidak puas	puas	
tahuB	tahu	Count	20	10	30
		% within tahuB	66,7%	33,3%	100,0%
Total		Count	20	10	30
		% within tahuB	66,7%	33,3%	100,0%

Chi-Square Tests

	Value
Pearson Chi-Square	.(a)
N of Valid Cases	30

a No statistics are computed because tahuB is a constant.

Risk Estimate

	Value
Odds Ratio for tahuB (tahu / .)	.(a)

a No statistics are computed because tahuB is a constant.