



Explore

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
umur	55	96,5%	2	3,5%	57	100,0%
lamapakai	55	96,5%	2	3,5%	57	100,0%
lamasemprotminggu	55	96,5%	2	3,5%	57	100,0%
terakhir	55	96,5%	2	3,5%	57	100,0%
rokok	55	96,5%	2	3,5%	57	100,0%
chol	55	96,5%	2	3,5%	57	100,0%

Tests of Normality

	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
umur	,102	55	,200*	,965	55	,111
lamapakai	,173	55	,000	,906	55	,000
lamasemprotminggu	,315	55	,000	,686	55	,000
terakhir	,275	55	,000	,749	55	,000
rokok	,197	55	,000	,837	55	,000
chol	,281	55	,000	,797	55	,000

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

Statistics

		umur	lamapakai	lamasempr otminggu	terakhir	chol	lamabertani
N	Valid	56	56	56	56	56	56
	Missing	1	1	1	1	1	1
Mean		41,2500	15,2321	7,0268	6,4107	48,6607	16,8750
Median		39,5000	11,0000	4,0000	4,0000	50,0000	15,0000
Mode		30,00	10,00	4,00	1,00 ^a	50,00	10,00 ^a
Std. Deviation		11,60133	11,78287	8,28151	7,47305	8,48748	12,61610
Variance		134,591	138,836	68,583	55,846	72,037	159,166
Minimum		18,00	1,00	,25	,00	37,50	1,00
Maximum		73,00	50,00	42,00	30,00	62,50	50,00

a. Multiple modes exist. The smallest value is shown

Descriptives

			Statistic	Std. Error	
umur	Mean		41,1273	1,57379	
	95% Confidence Interval for Mean	Lower Bound	37,9720		
		Upper Bound	44,2825		
	5% Trimmed Mean		40,7727		
	Median		39,0000		
	Variance		136,224		
	Std. Deviation		11,67151		
	Minimum		18,00		
	Maximum		73,00		
	Range		55,00		
	Interquartile Range		18,00		
	Skewness		,504		,322
	Kurtosis		-,019		,634
	lamapakai	Mean			14,9818
95% Confidence Interval for Mean		Lower Bound	11,8080		
		Upper Bound	18,1556		
5% Trimmed Mean			14,1667		
Median			10,0000		
Variance			137,833		
Std. Deviation			11,74023		
Minimum			1,00		
Maximum			50,00		
Range			49,00		
Interquartile Range			18,00		
Skewness			,956	,322	
Kurtosis			,369	,634	
lamasemprotminggu		Mean		7,0818	1,12558
	95% Confidence Interval for Mean	Lower Bound	4,8252		
		Upper Bound	9,3385		
	5% Trimmed Mean		6,0278		
	Median		4,0000		
	Variance		69,681		
	Std. Deviation		8,34750		
	Minimum		,25		
	Maximum		42,00		
	Range		41,75		
	Interquartile Range		4,00		
	Skewness		2,338	,322	
	Kurtosis		5,733	,634	
	terakhir	Mean		6,5273	
95% Confidence Interval for Mean		Lower Bound	4,5023		
		Upper Bound	8,5522		
5% Trimmed Mean			5,5859		
Median			4,0000		
Variance			56,106		
Std. Deviation			7,49038		
Minimum			,00		
Maximum			30,00		
Range			30,00		
Interquartile Range			6,00		
Skewness			1,902	,322	
Kurtosis			3,389	,634	
rokok		Mean		7,8545	1,00115
	95% Confidence Interval for Mean	Lower Bound	5,8474		
		Upper Bound	9,8617		
	5% Trimmed Mean		7,1111		
	Median		6,0000		
	Variance		55,127		
	Std. Deviation		7,42473		
	Minimum		,00		
	Maximum		32,00		
	Range		32,00		
	Interquartile Range		12,00		
	Skewness		1,108	,322	
	Kurtosis		1,828	,634	
	chol	Mean		48,8636	
95% Confidence Interval for Mean		Lower Bound	46,5854		
		Upper Bound	51,1419		
5% Trimmed Mean			48,7374		
Median			50,0000		
Variance			71,023		
Std. Deviation			8,42750		
Minimum			37,50		
Maximum			62,50		
Range			25,00		
Interquartile Range			12,50		
Skewness			,109	,322	
Kurtosis			-,729	,634	

Explore

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
terakhir	56	98,2%	1	1,8%	57	100,0%

Descriptives

			Statistic	Std. Error
terakhir	Mean		6,4107	,99863
	95% Confidence Interval for Mean	Lower Bound	4,4094	
		Upper Bound	8,4120	
	5% Trimmed Mean		5,4563	
	Median		4,0000	
	Variance		55,846	
	Std. Deviation		7,47305	
	Minimum		,00	
	Maximum		30,00	
	Range		30,00	
	Interquartile Range		6,00	
	Skewness		1,915	,319
	Kurtosis		3,459	,628

Frequency Table

choklat

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	1,00	40	70,2	71,4	71,4
	2,00	16	28,1	28,6	100,0
	Total	56	98,2	100,0	
Missing	System	1	1,8		
Total		57	100,0		

lama pakai pestisida

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	1,00	28	49,1	50,0	50,0
	2,00	28	49,1	50,0	100,0
	Total	56	98,2	100,0	
Missing	System	1	1,8		
Total		57	100,0		

lama semprot per minggu

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	1,00	32	56,1	57,1	57,1
	2,00	24	42,1	42,9	100,0
	Total	56	98,2	100,0	
Missing	System	1	1,8		
Total		57	100,0		

waktu terakhir menyemprot

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	1,00	31	54,4	56,4	56,4
	2,00	24	42,1	43,6	100,0
	Total	55	96,5	100,0	
Missing	System	2	3,5		
Total		57	100,0		

konsumsi rokok per hari

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	1,00	28	49,1	50,0	50,0
	2,00	28	49,1	50,0	100,0
	Total	56	98,2	100,0	
Missing	System	1	1,8		
Total		57	100,0		

Frequency Table

masker

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	1,00	13	22,8	23,2	23,2
	2,00	43	75,4	76,8	100,0
	Total	56	98,2	100,0	
Missing	System	1	1,8		
Total		57	100,0		

sarungtangan

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	1,00	5	8,8	8,9	8,9
	2,00	51	89,5	91,1	100,0
	Total	56	98,2	100,0	
Missing	System	1	1,8		
Total		57	100,0		

overall

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	1,00	45	78,9	80,4	80,4
	2,00	11	19,3	19,6	100,0
	Total	56	98,2	100,0	
Missing	System	1	1,8		
Total		57	100,0		

boot

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	1,00	27	47,4	48,2	48,2
	2,00	29	50,9	51,8	100,0
	Total	56	98,2	100,0	
Missing	System	1	1,8		
Total		57	100,0		

konsayur

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	1,00	56	98,2	100,0	100,0
Missing	System	1	1,8		
Total		57	100,0		

mandi

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	1,00	21	36,8	37,5	37,5
	2,00	35	61,4	62,5	100,0
	Total	56	98,2	100,0	
Missing	System	1	1,8		
Total		57	100,0		

cucitangan

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	1,00	56	98,2	100,0	100,0
Missing	System	1	1,8		
Total		57	100,0		

makanrokok

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	1,00	7	12,3	12,5	12,5
	2,00	49	86,0	87,5	100,0
	Total	56	98,2	100,0	
Missing	System	1	1,8		
Total		57	100,0		

penyu

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	1,00	18	31,6	32,1	32,1
	2,00	38	66,7	67,9	100,0
	Total	56	98,2	100,0	
Missing	System	1	1,8		
Total		57	100,0		

Statistics

waktu terakhir menyemprot

N	Valid	56
	Missing	1
Mean		1,4464
Median		1,0000
Mode		1,00
Std. Deviation		,50162
Variance		,252
Minimum		1,00
Maximum		2,00

waktu terakhir menyemprot

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	1,00	31	54,4	55,4	55,4
	2,00	25	43,9	44,6	100,0
	Total	56	98,2	100,0	
Missing	System	1	1,8		
Total		57	100,0		

masker * cholkat Crosstabulation

			cholkat		Total
			1,00	2,00	
masker	1,00	Count	11	2	13
		% within masker	84,6%	15,4%	100,0%
	2,00	Count	29	14	43
		% within masker	67,4%	32,6%	100,0%
Total		Count	40	16	56
		% within masker	71,4%	28,6%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1,443 ^b	1	,230		
Continuity Correction ^a	,724	1	,395		
Likelihood Ratio	1,577	1	,209		
Fisher's Exact Test				,308	,200
Linear-by-Linear Association	1,417	1	,234		
N of Valid Cases	56				

a. Computed only for a 2x2 table

b. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 3,71.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for masker (1,00 / 2,00)	2,655	,517	13,635
For cohort cholkat = 1,00	1,255	,919	1,713
For cohort cholkat = 2,00	,473	,123	1,814
N of Valid Cases	56		

sarungtangan * cholkat Crosstabulation

			cholkat		Total
			1,00	2,00	
sarungtangan	1,00	Count	4	1	5
		% within sarungtangan	80,0%	20,0%	100,0%
	2,00	Count	36	15	51
		% within sarungtangan	70,6%	29,4%	100,0%
Total		Count	40	16	56
		% within sarungtangan	71,4%	28,6%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,198 ^b	1	,657		
Continuity Correction ^a	,000	1	1,000		
Likelihood Ratio	,211	1	,646		
Fisher's Exact Test				1,000	,555
Linear-by-Linear Association	,194	1	,660		
N of Valid Cases	56				

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,43.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for sarung tangan (1,00 / 2,00)	1,667	,172	16,174
For cohort choklat = 1,00	1,133	,706	1,818
For cohort choklat = 2,00	,680	,112	4,130
N of Valid Cases	56		

overall * choklat Crosstabulation

			choklat		Total
			1,00	2,00	
overall	1,00	Count	32	13	45
		% within overall	71,1%	28,9%	100,0%
	2,00	Count	8	3	11
		% within overall	72,7%	27,3%	100,0%
Total		Count	40	16	56
		% within overall	71,4%	28,6%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,011 ^b	1	,915		
Continuity Correction ^a	,000	1	1,000		
Likelihood Ratio	,011	1	,915		
Fisher's Exact Test				1,000	,616
Linear-by-Linear Association	,011	1	,916		
N of Valid Cases	56				

a. Computed only for a 2x2 table

b. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 3,14.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for overall (1,00 / 2,00)	,923	,211	4,036
For cohort cholkat = 1,00	,978	,651	1,469
For cohort cholkat = 2,00	1,059	,364	3,083
N of Valid Cases	56		

boot * cholkat Crosstabulation

			cholkat		Total
			1,00	2,00	
boot	1,00	Count	18	9	27
		% within boot	66,7%	33,3%	100,0%
	2,00	Count	22	7	29
		% within boot	75,9%	24,1%	100,0%
Total		Count	40	16	56
		% within boot	71,4%	28,6%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,579 ^b	1	,447		
Continuity Correction ^a	,216	1	,642		
Likelihood Ratio	,580	1	,446		
Fisher's Exact Test				,558	,321
Linear-by-Linear Association	,569	1	,451		
N of Valid Cases	56				

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 7,71.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for boot (1,00 / 2,00)	,636	,198	2,046
For cohort cholkat = 1,00	,879	,628	1,230
For cohort cholkat = 2,00	1,381	,598	3,190
N of Valid Cases	56		

mandi * cholkat Crosstabulation

			cholkat		Total
			1,00	2,00	
mandi	1,00	Count	15	6	21
		% within mandi	71,4%	28,6%	100,0%
	2,00	Count	25	10	35
		% within mandi	71,4%	28,6%	100,0%
Total		Count	40	16	56
		% within mandi	71,4%	28,6%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,000 ^b	1	1,000		
Continuity Correction ^a	,000	1	1,000		
Likelihood Ratio	,000	1	1,000		
Fisher's Exact Test				1,000	,616
Linear-by-Linear Association	,000	1	1,000		
N of Valid Cases	56				

a. Computed only for a 2x2 table

b. 0 cells (.0%) have expected count less than 5. The minimum expected count is 6,00.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for mandi (1,00 / 2,00)	1,000	,302	3,312
For cohort choklat = 1,00	1,000	,710	1,408
For cohort choklat = 2,00	1,000	,425	2,352
N of Valid Cases	56		

makanrokok * choklat Crosstabulation

			choklat		Total
			1,00	2,00	
makanrokok	tidak	Count	36	13	49
		% within makanrokok	73,5%	26,5%	100,0%
ya	Count	4	3	7	
	% within makanrokok	57,1%	42,9%	100,0%	
Total	Count	40	16	56	
	% within makanrokok	71,4%	28,6%	100,0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,800 ^b	1	,371		
Continuity Correction ^a	,200	1	,655		
Likelihood Ratio	,749	1	,387		
Fisher's Exact Test				,395	,314
Linear-by-Linear Association	,786	1	,375		
N of Valid Cases	56				

a. Computed only for a 2x2 table

b. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 2,00.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for makanrokok (tidak / ya)	2,077	,409	10,555
For cohort cholkat = 1,00	1,286	,662	2,496
For cohort cholkat = 2,00	,619	,234	1,640
N of Valid Cases	56		

penyu * cholkat Crosstabulation

			cholkat		Total
			1,00	2,00	
penyu	1,00	Count	13	5	18
		% within penyu	72,2%	27,8%	100,0%
	2,00	Count	27	11	38
		% within penyu	71,1%	28,9%	100,0%
Total		Count	40	16	56
		% within penyu	71,4%	28,6%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,008 ^b	1	,928		
Continuity Correction ^a	,000	1	1,000		
Likelihood Ratio	,008	1	,928		
Fisher's Exact Test				1,000	,595
Linear-by-Linear Association	,008	1	,929		
N of Valid Cases	56				

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 5,14.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for penyuluhan (1,00 / 2,00)	1,059	,304	3,687
For cohort cholkat = 1,00	1,016	,716	1,444
For cohort cholkat = 2,00	,960	,392	2,351
N of Valid Cases	56		

lama pakai pestisida * cholkat Crosstabulation

			cholkat		Total
			1,00	2,00	
lama pakai pestisida	1,00	Count	22	6	28
		% within lama pakai pestisida	78,6%	21,4%	100,0%
	2,00	Count	18	10	28
		% within lama pakai pestisida	64,3%	35,7%	100,0%
Total		Count	40	16	56
		% within lama pakai pestisida	71,4%	28,6%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1,400 ^b	1	,237		
Continuity Correction ^a	,788	1	,375		
Likelihood Ratio	1,411	1	,235		
Fisher's Exact Test				,375	,188
Linear-by-Linear Association	1,375	1	,241		
N of Valid Cases	56				

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 8,00.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for lama pakai pestisida (1,00 / 2,00)	2,037	,621	6,686
For cohort cholkat = 1,00	1,222	,872	1,712
For cohort cholkat = 2,00	,600	,252	1,426
N of Valid Cases	56		

lama semprot per minggu * cholkat Crosstabulation

			cholkat		Total
			1,00	2,00	
lama semprot per minggu	1,00	Count	21	11	32
		% within lama semprot per minggu	65,6%	34,4%	100,0%
	2,00	Count	19	5	24
		% within lama semprot per minggu	79,2%	20,8%	100,0%
Total		Count	40	16	56
		% within lama semprot per minggu	71,4%	28,6%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1,232 ^b	1	,267		
Continuity Correction ^a	,658	1	,417		
Likelihood Ratio	1,259	1	,262		
Fisher's Exact Test				,373	,210
Linear-by-Linear Association	1,210	1	,271		
N of Valid Cases	56				

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 6,86.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for lama semprot per minggu (1,00 / 2,00)	,502	,147	1,712
For cohort cholkat = 1,00	,829	,600	1,146
For cohort cholkat = 2,00	1,650	,661	4,120
N of Valid Cases	56		

kebiasaan merokok y/n * cholkat Crosstabulation

			cholkat		Total
			1,00	2,00	
kebiasaan merokok y/n	tidak	Count	14	3	17
		% within kebiasaan merokok y/n	82,4%	17,6%	100,0%
	ya	Count	26	13	39
		% within kebiasaan merokok y/n	66,7%	33,3%	100,0%
Total	Count	40	16	56	
	% within kebiasaan merokok y/n	71,4%	28,6%	100,0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1,427 ^b	1	,232		
Continuity Correction ^a	,762	1	,383		
Likelihood Ratio	1,514	1	,219		
Fisher's Exact Test				,339	,193
Linear-by-Linear Association	1,402	1	,236		
N of Valid Cases	56				

a. Computed only for a 2x2 table

b. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 4,86.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for kebiasaan merokok y/n (tidak / ya)	2,333	,568	9,591
For cohort cholkat = 1,00	1,235	,904	1,688
For cohort cholkat = 2,00	,529	,173	1,620
N of Valid Cases	56		

waktu terakhir menyemprot * cholkat Crosstabulation

			cholkat		Total
			1,00	2,00	
waktu terakhir menyemprot	1,00	Count	24	7	31
		% within waktu terakhir menyemprot	77,4%	22,6%	100,0%
	2,00	Count	16	9	25
		% within waktu terakhir menyemprot	64,0%	36,0%	100,0%
Total		Count	40	16	56
		% within waktu terakhir menyemprot	71,4%	28,6%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1,221 ^b	1	,269		
Continuity Correction ^a	,652	1	,419		
Likelihood Ratio	1,217	1	,270		
Fisher's Exact Test				,374	,209
Linear-by-Linear Association	1,199	1	,273		
N of Valid Cases	56				

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 7,14.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for waktu terakhir menyemprot (1,00 / 2,00)	1,929	,597	6,232
For cohort cholkat = 1,00	1,210	,852	1,717
For cohort cholkat = 2,00	,627	,272	1,446
N of Valid Cases	56		



PETA LOKASI PENELITIAN

