

## LAMPIRAN 2

### Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
dekat laut * Hipertensi	1000	100,0%	0	,0%	1000	100,0%
murba * Hipertensi	1000	100,0%	0	,0%	1000	100,0%
Merokok * Hipertensi	1000	100,0%	0	,0%	1000	100,0%
Diabetes * Hipertensi	1000	100,0%	0	,0%	1000	100,0%
Obesitas * Hipertensi	1000	100,0%	0	,0%	1000	100,0%
Aktivitas Fisik * Hipertensi	1000	100,0%	0	,0%	1000	100,0%
Jenis Kelamin * Hipertensi	1000	100,0%	0	,0%	1000	100,0%
Tempat * Hipertensi	1000	100,0%	0	,0%	1000	100,0%
Poli Kunjungan * Hipertensi	1000	100,0%	0	,0%	1000	100,0%

### 1. Tempat \* Hipertensi

#### Crosstab

			Hipertensi		Total
			Ya	Tidak	Ya
Tempat	Penjaringan	Count	3	147	150
		% within Tempat	2,0%	98,0%	100,0%
	Kelapa Gading	Count	11	89	100
		% within Tempat	11,0%	89,0%	100,0%
	Tanjung Priok	Count	13	187	200
		% within Tempat	6,5%	93,5%	100,0%
	Cilincing	Count	33	217	250
		% within Tempat	13,2%	86,8%	100,0%
	Pademangan	Count	26	124	150
		% within Tempat	17,3%	82,7%	100,0%
	Koja	Count	33	117	150
		% within Tempat	22,0%	78,0%	100,0%
Total		Count	119	881	1000
		% within Tempat	11,9%	88,1%	100,0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	<b>38,885(a)</b>	<b>5</b>	<b>,000</b>
Likelihood Ratio	43,440	5	,000
Linear-by-Linear Association	34,287	1	,000
N of Valid Cases	1000		

a 0 cells (,0%) have expected count less than 5. The minimum expected count is 11,90.

### 1. Tanjung Priok VS Penjaringan

#### PKM Kecamatan \* tensi Crosstabulation

PKM Kecamatan			tensi		Total
			ya	tidak	ya
	Tanjung Priok	Count	13	187	200
		% within PKM Kecamatan	6,5%	93,5%	100,0%

	Penjaringan	Count	3	147	150
		% within PKM Kecamatan	2,0%	98,0%	100,0%
Total		Count	16	334	350
		% within PKM Kecamatan	4,6%	95,4%	100,0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	3,979(b)	1	,046		
<b>Continuity Correction(a)</b>	<b>3,014</b>	<b>1</b>	<b>,083</b>		
Likelihood Ratio	4,373	1	,037		
Fisher's Exact Test				,068	,038
Linear-by-Linear Association	3,967	1	,046		
N of Valid Cases	350				

a Computed only for a 2x2 table

b 0 cells (,0%) have expected count less than 5. The minimum expected count is 6,86.

#### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for PKM Kecamatan (Tanjung Priok / Penjaringan)	3,406	,953	12,176
<b>For cohort tensi = ya</b>	<b>3,250</b>	<b>,943</b>	<b>11,202</b>
For cohort tensi = tidak	,954	,914	,996
N of Valid Cases	350		

#### 2. Kelapa Gading VS Penjaringan

##### tempat \* tensi Crosstabulation

			tensi		Total
			ya	tidak	ya
tempat	kelapa gading	Count	11	89	100
		% within tempat	11,0%	89,0%	100,0%
	penjaringan	Count	3	147	150
		% within tempat	2,0%	98,0%	100,0%
Total		Count	14	236	250
		% within tempat	5,6%	94,4%	100,0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	9,193(b)	1	,002		
<b>Continuity Correction(a)</b>	<b>7,570</b>	<b>1</b>	<b>,006</b>		
Likelihood Ratio	9,193	1	,002		
Fisher's Exact Test				,004	,003
Linear-by-Linear Association	9,157	1	,002		
N of Valid Cases	250				

a Computed only for a 2x2 table

b 0 cells (,0%) have expected count less than 5. The minimum expected count is 5,60.

### Risk Estimate

	Value	95% CI	
	Lower	Upper	Lower
Odds Ratio for tempat (kelapa gading / penjaringa)	6,056	1,645	22,299
<b>For cohort tensi = ya</b>	<b>5,500</b>	<b>1,574</b>	<b>19,221</b>
For cohort tensi = tidak	,908	,845	,977
N of Valid Cases	250		

### 3. Cilincing VS Penjaringan

#### PKM Kecamatan \* tensi Crosstabulation

PKM Kecamatan			tensi		Total
			ya	tidak	ya
Cilincing	Count		33	217	250
	% within PKM Kecamatan		13,2%	86,8%	100,0%
Penjaringan	Count		3	147	150
	% within PKM Kecamatan		2,0%	98,0%	100,0%
Total		Count	36	364	400
		% within PKM Kecamatan	9,0%	91,0%	100,0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	14,359(b)	1	,000		
<b>Continuity Correction(a)</b>	<b>13,024</b>	<b>1</b>	<b>,000</b>		
Likelihood Ratio	17,533	1	,000		
Fisher's Exact Test				,000	,000
Linear-by-Linear Association	14,323	1	,000		
N of Valid Cases	400				

a Computed only for a 2x2 table

b 0 cells (,0%) have expected count less than 5. The minimum expected count is 13,50.

### Risk Estimate

	Value	95% CI	
	Lower	Upper	Lower
Odds Ratio for PKM Kecamatan (Cilincing / Penjaringan)	7,452	2,244	24,748
<b>For cohort tensi = ya</b>	<b>6,600</b>	<b>2,060</b>	<b>21,147</b>
For cohort tensi = tidak	,886	,840	,934
N of Valid Cases	400		

### 4. Pademangan VS Penjaringan

#### tempat \* tensi Crosstabulation

tempat			tensi		Total
			ya	tidak	ya
pademangan	Count		26	124	150
	% within tempat		17,3%	82,7%	100,0%
penjaringan	Count		3	147	150
	% within tempat		2,0%	98,0%	100,0%

Total	Count	29	271	300
	% within tempat	9,7%	90,3%	100,0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	20,193(b)	1	,000		
<b>Continuity Correction(a)</b>	<b>18,476</b>	<b>1</b>	<b>,000</b>		
Likelihood Ratio	22,866	1	,000		
Fisher's Exact Test				,000	,000
Linear-by-Linear Association	20,126	1	,000		
N of Valid Cases	300				

a Computed only for a 2x2 table

b 0 cells (,0%) have expected count less than 5. The minimum expected count is 14,50.

#### Risk Estimate

	Value	95% C I	
	Lower	Upper	Lower
Odds Ratio for tempat (pademangan / penjaringan)	10,274	3,037	34,757
<b>For cohort tensi = ya</b>	<b>8,667</b>	<b>2,681</b>	<b>28,021</b>
For cohort tensi = tidak	,844	,781	,911
N of Valid Cases	300		

#### 5. Koja VS Penjaringan

##### tempat \* tensi Crosstabulation

tempat			tensi		Total ya
			ya	TIDAK	
koja	Count	33	117	150	
	% within tempat	22,0%	78,0%	100,0%	
penjaringan	Count	3	147	150	
	% within tempat	2,0%	98,0%	100,0%	
Total		Count	36	264	300
		% within tempat	12,0%	88,0%	100,0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	28,409(b)	1	,000		
<b>Continuity Correction(a)</b>	<b>26,547</b>	<b>1</b>	<b>,000</b>		
Likelihood Ratio	32,671	1	,000		
Fisher's Exact Test				,000	,000
Linear-by-Linear Association	28,314	1	,000		
N of Valid Cases	300				

a Computed only for a 2x2 table

b 0 cells (,0%) have expected count less than 5. The minimum expected count is 18,00.

#### Risk Estimate

	Value	95% C I	
	Lower	Upper	Lower

Odds Ratio for tempat (koja / penjaringan)	13,821	4,135	46,189
<b>For cohort tensi = ya</b>	<b>11,000</b>	<b>3,448</b>	<b>35,090</b>
For cohort tensi = TIDAK	,796	,729	,869
N of Valid Cases	300		

## 2. Poli Kunjungan \* Hipertensi

### Crosstab

Poli Kunjungan			Hipertensi		Total
			Ya	Tidak	Ya
BP Askes	Count		10	129	139
	% within Poli Kunjungan		7,2%	92,8%	100,0%
BP Umum	Count		53	666	719
	% within Poli Kunjungan		7,4%	92,6%	100,0%
BP Lansia	Count		56	86	142
	% within Poli Kunjungan		39,4%	60,6%	100,0%
Total		Count	119	881	1000
		% within Poli Kunjungan	11,9%	88,1%	100,0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	<b>119,705(a)</b>	<b>2</b>	<b>,000</b>
Likelihood Ratio	89,092	2	,000
Linear-by-Linear Association	70,648	1	,000
N of Valid Cases	1000		

a 0 cells (.0%) have expected count less than 5. The minimum expected count is 16,54

### 1. Umum VS Asuransi

#### yankes baru \* tensi Crosstabulation

yankes baru			tensi		Total
			ya	tidak	ya
BP UMUM	Count		53	666	719
	% within yankes baru		7,4%	92,6%	100,0%
BP Asuransi	Count		10	129	139
	% within yankes baru		7,2%	92,8%	100,0%
Total		Count	63	795	858
		% within yankes baru	7,3%	92,7%	100,0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,005(b)	1	,942		
<b>Continuity Correction(a)</b>	<b>,000</b>	<b>1</b>	<b>1,000</b>		
Likelihood Ratio	,005	1	,941		
Fisher's Exact Test				1,000	,555
Linear-by-Linear Association	,005	1	,942		
N of Valid Cases	858				

a Computed only for a 2x2 table

b 0 cells (.0%) have expected count less than 5. The minimum expected count is 10,21.

### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for yankes baru (BP UMUm / BP Asuransi)	1,027	,509	2,070
<b>For cohort tensi = ya</b>	<b>1,025</b>	<b>,534</b>	<b>1,964</b>
For cohort tensi = tidak	,998	,949	1,050
N of Valid Cases	858		

### 2. Lansia VS Asuransi

#### yankes baru \* tensi Crosstabulation

yankes baru			tensi		Total
			ya	tidak	ya
BP lansia	Count		56	86	142
	% within yankes baru		39,4%	60,6%	100,0%
BP Asuransi	Count		10	129	139
	% within yankes baru		7,2%	92,8%	100,0%
Total		Count	66	215	281
		% within yankes baru	23,5%	76,5%	100,0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	40,633(b)	1	,000		
<b>Continuity Correction(a)</b>	<b>38,859</b>	<b>1</b>	<b>,000</b>		
Likelihood Ratio	43,978	1	,000		
Fisher's Exact Test				,000	,000
Linear-by-Linear Association	40,489	1	,000		
N of Valid Cases	281				

a Computed only for a 2x2 table

b 0 cells (,0%) have expected count less than 5. The minimum expected count is 32,65.

### Risk Estimate

	Value	95% C I	
		Lower	Upper
Odds Ratio for yankes baru (BP lansia / BP Asuransi)	8,400	4,064	17,362
<b>For cohort tensi = ya</b>	<b>5,482</b>	<b>2,917</b>	<b>10,302</b>
For cohort tensi = tidak	,653	,567	,751
N of Valid Cases	281		

### 3. Daerah pantai \* Hipertensi

#### Crosstab

			Hipertensi		Total
			Ya	Tidak	Ya
dekat laut	daerah pantai	Count	75	675	750
		% within dekat laut	10,0%	90,0%	100,0%
	bukan daerah pantai	Count	44	206	250

		% within dekat laut	17,6%	82,4%	100,0%
Total		Count	119	881	1000
		% within dekat laut	11,9%	88,1%	100,0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	10,330(b)	1	,001		
<b>Continuity Correction(a)</b>	<b>9,618</b>	<b>1</b>	<b>,002</b>		
Likelihood Ratio	9,594	1	,002		
Fisher's Exact Test				,002	,001
Linear-by-Linear Association	10,320	1	,001		
N of Valid Cases	1000				

a Computed only for a 2x2 table

b 0 cells (,0%) have expected count less than 5. The minimum expected count is 29,75.

#### Risk Estimate

	Value	95% C I	
	Lower	Upper	Lower
Odds Ratio for dekat laut (daerah pantai / bukan daerah pantai)	,520	,347	,779
<b>For cohort Hipertensi = Ya</b>	<b>,568</b>	<b>,403</b>	<b>,801</b>
For cohort Hipertensi = Tidak	1,092	1,027	1,162
N of Valid Cases	1000		

#### 4. Umur baru \* Hipertensi

##### Crosstab

murba			Hipertensi		Total
			Ya	Tidak	Ya
25-39 tahun	Count	6	424	430	
	% within murba	1,4%	98,6%	100,0%	
40-44 tahun	Count	7	71	78	
	% within murba	9,0%	91,0%	100,0%	
45-49 tahun	Count	8	102	110	
	% within murba	7,3%	92,7%	100,0%	
50-54 tahun	Count	19	110	129	
	% within murba	14,7%	85,3%	100,0%	
55-59 tahun	Count	42	116	158	
	% within murba	26,6%	73,4%	100,0%	
60-64 tahun	Count	37	58	95	
	% within murba	38,9%	61,1%	100,0%	
Total		Count	119	881	1000
		% within murba	11,9%	88,1%	100,0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	<b>147,906(a)</b>	<b>5</b>	<b>,000</b>
Likelihood Ratio	144,395	5	,000

Linear-by-Linear Association	128,723	1	,000
N of Valid Cases	1000		

a 0 cells (,0%) have expected count less than 5. The minimum expected count is 9,28.

## Crosstabs

### 1. 40-44 vs 25-39

#### murba \* tensi Crosstabulation

		tensi		Total	
		ya	tidak	ya	
murba	40-44	Count	7	71	78
		% within murba	9,0%	91,0%	100,0%
	25-39	Count	6	423	429
		% within murba	1,4%	98,6%	100,0%
Total		Count	13	494	507
		% within murba	2,6%	97,4%	100,0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	15,161(b)	1	,000		
<b>Continuity Correction(a)</b>	<b>12,281</b>	<b>1</b>	<b>,000</b>		
Likelihood Ratio	10,661	1	,001		
Fisher's Exact Test				,001	,001
Linear-by-Linear Association	15,132	1	,000		
N of Valid Cases	507				

a Computed only for a 2x2 table

b 1 cells (25,0%) have expected count less than 5. The minimum expected count is 2,00.

#### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for murba (40-44 / 25-39)	6,951	2,270	21,282
<b>For cohort tensi = ya</b>	<b>6,417</b>	<b>2,216</b>	<b>18,584</b>
For cohort tensi = tidak	,923	,860	,991
N of Valid Cases	507		

### 2. 45-49 vs 25-39

#### murba \* tensi Crosstabulation

		tensi		Total	
		ya	tidak	ya	
murba	45-49	Count	8	102	110
		% within murba	7,3%	92,7%	100,0%
	25-39	Count	6	424	430
		% within murba	1,4%	98,6%	100,0%
Total		Count	14	526	540
		% within murba	2,6%	97,4%	100,0%

#### Chi-Square Tests



	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	11,981(b)	1	,001		
<b>Continuity Correction(a)</b>	<b>9,767</b>	<b>1</b>	<b>,002</b>		
Likelihood Ratio	9,384	1	,002		
Fisher's Exact Test				,002	,002
Linear-by-Linear Association	11,959	1	,001		
N of Valid Cases	540				

a Computed only for a 2x2 table

b 1 cells (25,0%) have expected count less than 5. The minimum expected count is 2,85.

#### Risk Estimate

	Value	95% C I	
	Lower	Upper	Lower
Odds Ratio for murba (45-49 / 25-39)	5,542	1,882	16,326
<b>For cohort tensi = ya</b>	<b>5,212</b>	<b>1,847</b>	<b>14,711</b>
For cohort tensi = tidak	,940	,891	,992
N of Valid Cases	540		

### 3. 50-54 vs 25-39

#### murba \* tensi Crosstabulation

			tensi		Total
			ya	tidak	ya
murba	50-54	Count	19	110	129
		% within murba	14,7%	85,3%	100,0%
	25-39	Count	6	424	430
		% within murba	1,4%	98,6%	100,0%
Total		Count	25	534	559
		% within murba	4,5%	95,5%	100,0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	41,292(b)	1	,000		
<b>Continuity Correction(a)</b>	<b>38,230</b>	<b>1</b>	<b>,000</b>		
Likelihood Ratio	33,211	1	,000		
Fisher's Exact Test				,000	,000
Linear-by-Linear Association	41,218	1	,000		
N of Valid Cases	559				

a Computed only for a 2x2 table

b 0 cells (,0%) have expected count less than 5. The minimum expected count is 5,77.

#### Risk Estimate

	Value	95% C I	
	Lower	Upper	Lower
Odds Ratio for murba (50-54 / 25-39)	12,206	4,761	31,294
<b>For cohort tensi = ya</b>	<b>10,556</b>	<b>4,307</b>	<b>25,872</b>

For cohort tensi = tidak	,865	,804	,930
N of Valid Cases	559		

#### 4. 55-59 vs 25-39

##### murba \* tensi Crosstabulation

		tensi		Total	
		ya	tidak	ya	
murba	55-59	Count	42	116	158
		% within murba	26,6%	73,4%	100,0%
	25-39	Count	6	424	430
		% within murba	1,4%	98,6%	100,0%
Total		Count	48	540	588
		% within murba	8,2%	91,8%	100,0%

##### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	97,773(b)	1	,000		
<b>Continuity Correction(a)</b>	<b>94,442</b>	<b>1</b>	<b>,000</b>		
Likelihood Ratio	86,338	1	,000		
Fisher's Exact Test				,000	,000
Linear-by-Linear Association	97,607	1	,000		
N of Valid Cases	588				

a Computed only for a 2x2 table

b 0 cells (,0%) have expected count less than 5. The minimum expected count is 12,90.

##### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for murba (55-59 / 25-39)	25,586	10,616	61,668
<b>For cohort tensi = ya</b>	<b>19,051</b>	<b>8,259</b>	<b>43,941</b>
For cohort tensi = tidak	,745	,677	,818
N of Valid Cases	588		

#### 5. 60-64 vs 25-39

##### murba \* tensi Crosstabulation

		tensi		Total
		ya	tidak	ya
murba	60-64	37	58	95
	25-39	6	424	430
Total		43	482	525

##### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	145,916(b)	1	,000		
<b>Continuity Correction(a)</b>	<b>140,965</b>	<b>1</b>	<b>,000</b>		

Likelihood Ratio	107,369	1	,000		
Fisher's Exact Test				,000	,000
Linear-by-Linear Association	145,638	1	,000		
N of Valid Cases	525				

a Computed only for a 2x2 table

b 0 cells (,0%) have expected count less than 5. The minimum expected count is 7,78.

#### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for murba (60-64 / 25-39)	45,080	18,234	111,456
<b>For cohort tensi = ya</b>	<b>27,912</b>	<b>12,129</b>	<b>64,235</b>
For cohort tensi = tidak	,619	,527	,727
N of Valid Cases	525		

#### 5. Jenis Kelamin \* Hipertensi

##### Crosstab

			Hipertensi		Total
			Ya	Tidak	Ya
Jenis Kelamin	Laki-Laki	Count	52	381	433
		% within Jenis Kelamin	12,0%	88,0%	100,0%
	Perempuan	Count	67	500	567
		% within Jenis Kelamin	11,8%	88,2%	100,0%
Total		Count	119	881	1000
		% within Jenis Kelamin	11,9%	88,1%	100,0%

##### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,009(b)	1	,926		
<b>Continuity Correction(a)</b>	<b>,000</b>	<b>1</b>	<b>1,000</b>		
Likelihood Ratio	,009	1	,926		
Fisher's Exact Test				,922	,501
Linear-by-Linear Association	,009	1	,926		
N of Valid Cases	1000				

a Computed only for a 2x2 table

b 0 cells (,0%) have expected count less than 5. The minimum expected count is 51,53.

#### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Jenis Kelamin (Laki-Laki / Perempuan)	1,019	,692	1,498
<b>For cohort Hipertensi = Ya</b>	<b>1,016</b>	<b>,723</b>	<b>1,428</b>
For cohort Hipertensi = Tidak	,998	,953	1,045
N of Valid Cases	1000		

## 6. Diabetes \* Hipertensi

### Crosstab

			Hipertensi		Total
			Ya	Tidak	Ya
Diabetes	Ya	Count	27	82	109
		% within Diabetes	24,8%	75,2%	100,0%
	Tidak	Count	92	799	891
		% within Diabetes	10,3%	89,7%	100,0%
Total		Count	119	881	1000
		% within Diabetes	11,9%	88,1%	100,0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	19,330(b)	1	,000		
<b>Continuity Correction(a)</b>	<b>17,976</b>	<b>1</b>	<b>,000</b>		
Likelihood Ratio	15,881	1	,000		
Fisher's Exact Test				,000	,000
Linear-by-Linear Association	19,310	1	,000		
N of Valid Cases	1000				

a Computed only for a 2x2 table

b 0 cells (,0%) have expected count less than 5. The minimum expected count is 12,97.

### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Diabetes (Ya / Tidak)	2,860	1,760	4,647
<b>For cohort Hipertensi = Ya</b>	<b>2,399</b>	<b>1,640</b>	<b>3,508</b>
For cohort Hipertensi = Tidak	,839	,752	,936
N of Valid Cases	1000		

## 7. Obesitas \* Hipertensi

### Crosstab

			Hipertensi		Total
			Ya	Tidak	Ya
Obesitas	Ya	Count	11	50	61
		% within Obesitas	18,0%	82,0%	100,0%
	Tidak	Count	108	831	939
		% within Obesitas	11,5%	88,5%	100,0%
Total		Count	119	881	1000
		% within Obesitas	11,9%	88,1%	100,0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	2,331(b)	1	,127		

<b>Continuity Correction(a)</b>	<b>1,749</b>	<b>1</b>	<b>,186</b>		
Likelihood Ratio	2,073	1	,150		
Fisher's Exact Test				,150	,097
Linear-by-Linear Association	2,328	1	,127		
N of Valid Cases	1000				

a Computed only for a 2x2 table

b 0 cells (,0%) have expected count less than 5. The minimum expected count is 7,26.

#### Risk Estimate

	Value	95% Confidence Interval	
	Lower	Upper	Lower
Odds Ratio for Obesitas (Ya / Tidak)	1,693	,855	3,351
<b>For cohort Hipertensi = Ya</b>	<b>1,568</b>	<b>,892</b>	<b>2,755</b>
For cohort Hipertensi = Tidak	,926	,822	1,044
N of Valid Cases	1000		

#### 8. Aktivitas Fisik \* Hipertensi

##### Crosstab

		Hipertensi		Total	
		Ya	Tidak	Ya	
Aktivitas Fisik	Tidak	Count	16	61	77
		% within Aktivitas Fisik	20,8%	79,2%	100,0%
	Ya	Count	103	820	923
		% within Aktivitas Fisik	11,2%	88,8%	100,0%
Total		Count	119	881	1000
		% within Aktivitas Fisik	11,9%	88,1%	100,0%

##### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	6,274(b)	1	,012		
<b>Continuity Correction(a)</b>	<b>5,390</b>	<b>1</b>	<b>,020</b>		
Likelihood Ratio	5,369	1	,021		
Fisher's Exact Test				,017	,014
Linear-by-Linear Association	6,267	1	,012		
N of Valid Cases	1000				

a Computed only for a 2x2 table

b 0 cells (,0%) have expected count less than 5. The minimum expected count is 9,16.

#### Risk Estimate

	Value	95% C I	
	Lower	Upper	Lower
Odds Ratio for Aktivitas Fisik (Tidak / Ya)	2,088	1,161	3,757
<b>For cohort Hipertensi = Ya</b>	<b>1,862</b>	<b>1,161</b>	<b>2,987</b>
For cohort Hipertensi = Tidak	,892	,794	1,002
N of Valid Cases	1000		

## 9. Merokok \* Hipertensi

### Crosstab

			Hipertensi		Total
			Ya	Tidak	Ya
Merokok	Ya	Count	22	242	264
		% within Merokok	8,3%	91,7%	100,0%
	tidak	Count	97	639	736
		% within Merokok	13,2%	86,8%	100,0%
Total		Count	119	881	1000
		% within Merokok	11,9%	88,1%	100,0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	4,352(b)	1	,037		
<b>Continuity Correction(a)</b>	<b>3,902</b>	<b>1</b>	<b>,048</b>		
Likelihood Ratio	4,647	1	,031		
Fisher's Exact Test				,036	,022
Linear-by-Linear Association	4,348	1	,037		
N of Valid Cases	1000				

a Computed only for a 2x2 table

b 0 cells (.0%) have expected count less than 5. The minimum expected count is 31,42.

### Risk Estimate

	Value	95% Confidence Interval	
	Lower	Upper	Lower
Odds Ratio for Merokok (Ya / tidak)	,599	,368	,974
<b>For cohort Hipertensi = Ya</b>	<b>,632</b>	<b>,407</b>	<b>,983</b>
For cohort Hipertensi = Tidak	1,056	1,008	1,106
N of Valid Cases	1000		